

1. Alonso P, García J, Gómez M et al.: Rentability of anaesthetic procedure. 8th European Congress Anaesthesiology 1990. 5; 6; 2-6.
2. Alonso, J; Prieto, L; Antó JM.: La versión española del SF-36 health survey: un instrumento para la medida de los resultados clínicos. *Medicina Clínica*, 104:771-776, 1995.
3. Amarigiri SV, Lees TA. Elastic compression stockings for prevention of deep vein thrombosis. *Cochrane Review*. The Cochrane Library, Issue 1, 2001. Oxford: Update Software.
4. Bellamy, N; Watson, W; Goldsmith, CH; Campbell, J; Stitt, LW: Validation study of WOMAC: A health status instrument for measuring clinically important patient relevant outcomes to antirheumatic drug therapy in patients with osteoarthritis of the hip or knee. *The Journal of Rheumatology* 1988; 15(12):1833-1840.
5. Bergqvist D, Jonsson B. Cost-effectiveness of prolonged out-of-hospital prophylaxis with low-molecular-weight heparin following total hip replacement. *Haemostasis* 2000;30 Suppl 2:130-5.
6. Bick RL, Haas SK. International Consensus Recommendations. Summary statement and additional suggested guidelines. *Med Clin N Am* 1998; 82(3): 613-633.
7. Bleri C, Sztan M., Fourgeaux B et al.: Evaluation of a protocol for selective ordering of preoperative tests. *Lancet* 1986 1. 139-141
8. Callaghan J.J., Dennis D.A., Paprosky W.G., Rosenberg A.G.: Orthopaedic Knowledge Update. Hip and Knee reconstruction. Publicado por la AAOS.1995.Cap.22-23:179-206.
9. Circular 10/2001, de 11 de Octubre, del Ministerio de Sanidad y Consumo (BOE núm. 265, de lunes 5 de noviembre de 2001, pág. 40.133).
10. Clagett GP, Anderson FA, Geerts W, et al. Prevention of venous thromboembolism. Fifth ACCP Consensus Conference on Antithrombotic Therapy. *Chest* 1998; 114: 531S-560S.
11. Covert CR, Fox GS. Anaesthesia for hip surgery in the elderly. *Can J Anaesth* 1989 May;36(3 Pt 1):311-9.
12. Davis FM, Woolner DF, Frampton C, Wilkinson A, Grant A, Harrison RT, Roberts MT.
13. Diagnóstico de enfermería. L. J. Carpenito. Interamericana. 3ª edición 1991.
14. Diagnósticos enfermeros de la NANDA. Definiciones y clasificación 1997-1998. Harcourt Brace.

15. Diagnósticos enfermeros. M.T. Luis. Mosby-Doyma Libros. Madrid 1996.
16. E. Contreras Fernández; A. Báez Cabeza; y otros. Planes de Cuidados Enfermeros Estandarizados En Atención Primaria. Ed. Distrito Sanitario Costa del Sol. SAS.
17. Espehaug B, Havelin LI, Engesaeter LB, Langeland N, Vollset SE. Patient-related risk factors for early revision of total hip replacements. A population register-based case-control study of 674 revised hips. *Acta Orthop Scand* 1997; 68(3): 207-15.
18. Espinosa Almendro, J.M.; Muñoz Cobos, F; y otros (1997): Atención al Mayor en Medicina Familiar. Ed. SAMFyC. Granada.
19. Faulker A, Kennedy LG, Baxter K, Donovan J, Wilkinson M, Bevan G: Effectiveness of hip prostheses in primary total hip replacement: a critical review of evidence and economic model. *Health Technology Assessment* 1998, 2(6): 1-133.
20. Freedman KB, Brookenthal KR, Fitzgerald RH, Williams S, Lonner JH.. A meta-analysis of thromboembolic prophylaxis following elective total hip arthroplasty. *JBJS-Am* 2000; 82: 929-38.
21. Glenny AM, Song F: Antimicrobial prophylaxis in total hip replacement: a systematic review. *Health Technol Assess* 1999, 3(21): 1-57.
22. Harris, WH.: Traumatic arthritis of the hip after dislocation and acetabular fractures: treatment by mold arthroplasty. *J Bone Joint Surg (Am)* 1969; 51-A: 737-55.
23. Henderson, V (1996); Virginia Henderson. Los primeros noventa años. Ed. Masson, S.A. Barcelona.
24. Hozack, WJ; Rothman, RH; Albert, TJ; Balderston, RA; Engh K.: Relationship of total hip arthroplasty outcomes to other orthopaedic procedures. *Clin Orthop*, 344:88-93, 1997.
25. Hull RD, Pineo GF, Stein PD et al. Extended out-of-hospital low-molecular-weight heparin prophylaxis against deep venous thrombosis in patients after elective hip arthroplasty: a systematic review. *Ann Intern Med* 2001; 135: 858-69.
26. Hull RD, Pineo GF. Prophylaxis of deep venous thrombosis and pulmonary embolism. Current recommendations. *Med Clin N Am* 1998; 82(3): 477-93.
27. Hull RD. New insights into extended prophylaxis after orthopaedic surgery - the North American Fragmin Trial experience. *Haemostasis* 2000;30 Suppl 2:95-100.
28. Kaplan EB, Boeckann AS, Roizen MF et al.: Elimination of unnecessary preoperative laboratory tests, *Anesthesiology* 1982, 57; A 445.
29. Koval KJ, Aharonoff GB, Rosenberg AD, Schmigelski C, Bernstein RL, Zuckerman JD. Hip fracture in the elderly: the effect of anaesthetic technique. *Orthopaedics* 1999 Jan;22(1):31-4.
30. Laberge A, Bernard PM, Lamarche PA. *Rev Epidemiol Sante Publique* 1997;45(1):5-12.
31. Luis Rodrigo, M.T.; Fernández Ferrín, C; y Navarro Gómez Mª V.(1998): De la teoría a la práctica. El pensamiento de V. Henderson en el siglo XXI. Ed. Masson, S.A. Barcelona.

32. McKenzie PJ, Wishart HY, Gray I, Smith G. *BR J Anaesth* 1985 Sep;57(9):853-7.
33. New York Heart Association Classification of Heart Diseases. New York: McGill Ed. 1997.
34. O'Hara DA, Duff A, Berlin JA, Poses RM, Lawrence VA, Huber EC, Noveck H, Strom BL, Carson JL. The effect of anaesthetic technique on postoperative outcomes in hip fracture repair. *Anesthesiology*. 2000 Apr;92(4):928-30.
35. Parker M.J., Rajan D.: Arthroplasties (with and Without bone cement) for proximal femoral fractures in adults (Cochrane Review). In *Cochrane Library*, Issue 3, 2001.
36. Parker MJ, Urwin SC, Handoll HH, Griffiths R. General versus spinal/epidural anaesthesia for surgery for hip fractures in adults (Cochrane Review). In: *The Cochrane Library*, Issue 1, 2001. Oxford: Update Software.
37. Poon PC, Rennie J, Gray DH. Review of total hip replacement. The Middlemore Hospital experience, 1980-1991. *N Z Med J* 2001; 114 (1133):254-6.
38. Prevention of venous thromboembolism. International Consensus Statement (Guidelines according to scientific evidence). *Int Angiol* 1997; 16: 3-38.
39. Quintana J.M., Escobar A., Goenaga J.I., Arostegui I., Beldarrain I. y Letona J.: Análisis de la evidencia científica sobre la efectividad y eficiencia de los diferentes tipos de artroplastia de cadera. Recomendaciones. Investigación comisionada. Vitoria-Gastéiz. Departamento de Sanidad, Gobierno Vasco 2000. Informe nº: Osteba D-00-07.
40. Racle JP, Benkhadra A, Poy JY, Gleizal B, Gaudray A. Comparative study of general and spinal anaesthesia in elderly women in hip surgery. *Ann Fr Anesth Reanim* 1986;5(1):24-30.
41. Riordan P, Bickler G, Lyons C. Lessons of a hip failure. Registers of joint replacement operations should be set up. *BMJ* 1998; 316(7149):1985.
42. Ritter, MA; Albohm, MJ.: Overview: Maintaining outcomes for total hip arthroplasty: The past, present and future. *Clin Orthop*. 344:81-84, 1997.
43. Rodgers A., Walker N., Schug S., McKee A. & cols.: Reduction on postoperative mortality and morbidity with epidural or spinal anaesthesia: results from overview of randomised trials. *Br Med J* 2000. 321 (7275) 1493-7.]
44. Roizen H.: Preoperative patient evaluation. *Canadian J Anaesth* 1989. 30-7.
45. Scottish Intercollegiate Guidelines Network. Management of elderly people with fractured hip. A national Clinical Guideline recommended for use in Scotland. July 1997.
46. Serrano P., López J., Duque B., et al. Patrón de utilización de pruebas preoperatorias en una población sana y asintomática en Canarias. *Rev. Esp. Anestesiología. Reanim.* 2001; 48: 307-313.
47. SF-36 Health survey. Copyright 19924 Medical Outcomes Trust. Reproducido con permiso.
48. Sorenson RM, Pace NL. *Anesthesiology* 1992 Dec;77 (6):1095-104.

49. Thadaka R; Br J Anaesth 1987;59 (9):1080-8.
50. Valentin N, Lomholt B, Jensen JS, Hejgaard N, Kreiner S. Spinal or general anaesthesia for surgery of the fractured hip? A prospective study of mortality in 578 patients. Br J Anaesth 1986 Mar;58(3):284-91.
51. Wilmore D., Kehlet H.: Management of patients in fast track surgery. Br Med J 2001. 322 (7284) 473-6.