

Adams N, Sim J. An overview of fibromyalgia syndrome: mechanisms, differential diagnosis and treatment approaches. *Physiotherapy* 1998; 84: 304-318.

Alonso Álvarez B. Ejercicio físico en la fibromialgia. *Rehabilitación* 2003; 37(6): 363-74.

Ansari A. The efficacy of newer antidepressants in the treatment of chronic pain: a review of current literature. *Harv Rev Psychiatry* 2000; 7: 257-77.

Ballina García FJ. En Manual SER de las Enfermedades Reumáticas, Capítulo 23, 112-115, Editorial Médica Panamericana S.A. 2004.

Bennet RM, Burckhardt CS, Clark SR, O'Reilly CA, Wiens AN, Campbell SM. Group Treatment of Fibromyalgia: A 6 Month Outpatient Program. *J Rheum* 1995; 23: 521-8.

Bennett RM, Gatter RA, Campbell SM, et al. A comparison of cyclobenzaprine and placebo in the management of fibrositis. *Arthritis Rheum*. 1988; 31: 1535-1542.

Bennett RM, Kamin M, Karin R, Rosenthal N. Tramadol and acetaminophen combination tablets in the treatment of fibromyalgia pain. *Am J Med* 2003; 114: 537-545.

Biasi G, Manca S, Marcolongo R. Tramadol in the fibromyalgia syndrome: a controlled clinical trial versus placebo. *Int J Clin Pharmacol Res*. 1988; 18: 13-19.

Blechman EA, Brownell KD. *Medicina conductual de la mujer*. Ediciones Martínez Roca SA; 1992.

Bosch Romero E, Saenz Moya N, Valls Esteve M, Vinolas Valer S. Estudio de la calidad de vida en pacientes con fibromialgia: impacto de un programa de educación sanitaria. *Atencion Primaria*. 2002; 30: 16-21.

Burckhardt CS, Clark SR, Bennet RM. The Fibromyalgia Impact Questionnaire: Development and Validation. *J Rheum* 1991; 18: 728-33.

Busch A, Schachter CL, Peloso PM, et al. Exercise for treating fibromyalgia syndrome (Cochrane Review). In *The Cochrane Library*, Issue 3; 2002.

Buskila D, Neumann L. Assessing Functional Disability and Health Status of Women with Fibromyalgia: Validation of a Hebrew Version of the Fibromyalgia Impact Questionnaire. *J Rheum* 1996; 23: 903-6.

Calfas KJ, Long BJ, Sallis JF, et al. A controlled trial of provider counseling to promote the adoption of physical activity. *Prev Med* 1996; 25: 225-233.

Campos Sánchez S, Epidemiología de la fibromialgia. *Rev Esp Reumatol* 2000; 27: 417-420.

Carette S, Bell MJ, Reynolds WJ, et al. Comparison of amitriptyline, cyclobenzaprine and placebo in the treatment of fibromyalgia; a randomized, double-blind clinical trial. *Arthritis Rheum.* 1994; 37: 32-40.

Carette S, McCain GA, Bell DA, Fam AG. Evaluation of amitriptyline in primary fibrositis. *Arthritis Rheum.* 1986; 29: 655-659.

Collado A, Alijotas J, Benito P et al. Documento de consenso sobre el diagnóstico y tratamiento de la fibromialgia en Cataluña. *Med Clin (Barc)* 2002; 118(19): 745-9.

Crofford L, Rusell IJ, Mease P, Corbing A, Young J, Lamoreaux L et al. Pregabalin improves pain associated with fibromyalgia syndrome in a multicenter, randomized placebo controlled monotherapy trial. *Arthritis Rheum* 2002; 46 Suppl: S613.

Dawson KA, Tiidus PM, Pierrynowski M, et al. Evaluation of a community-based exercise program for diminishing symptoms of fibromyalgia. *Physiother Can* 2003, 55: 17-22.

Davis M, McKay M, Robbins E. Técnicas de autocontrol emocional. Ediciones Martínez Roca SA 2001.

Fernández Rodríguez M.A. Fibromialgia. *Rev Esp Rheumatol* 2000; 27: 442-446.

Ferracioli G, Chirelli L, Scita F, et al. EMG-biofeedback training in fibromyalgia syndrome. *J Rheumatol.* 1987; 14: 820-5.

Fulcher KY; White PD. Randomised controlled trial of graded exercise in patients with the chronic fatigue syndrome. *BMJ* 1997; 314: 1647-1652.

García López A; Campos Sánchez S. Impacto socioeconómico de la fibromialgia. *Rev Esp Reumatol* 2000; 27: 447-449.

Geel SE, Robergs RA. The effect of graded resistance exercise on fibromyalgia symptoms and muscle bioenergetics: A pilot study. *Arthritis Care Res* 2002, 47: 82-86.

Glass JM; Lyden AK; Petzke F; Stein P; Whelen G; Ambrose K; Chrousos G; Clauw DJ. The effect of brief exercise cessation on pain, fatigue, and mood symptom development in healthy, fit individuals. *J Psychosom Res* 2004; 57(4): 391-398.

Godfrey RG. A guide to the understanding and use of tricyclic antidepressants in the overall management of fibromyalgia and other chronic pain syndromes. *Arch Intern Med* 1996; 156: 1047-52.

Goldenberg D, Mayskiy M, Mossey C, Ruthazer R, Smichd C. A randomized, double blind crossover trial of fluoxetine and amitriptyline in the treatment of fibromyalgia. *Arthritis Rheum* 1996; 39: 1852-9.

Goldenberg DL, Burckhardt C, Crofford L. Management of Fibromyalgia Syndrome. *Jama* 2004; 292: 2388-95.

Goldenberg DL, Felson DT, Dinerman HA. Randomized, controlled trial of amitriptyline and naproxen in the treatment of patients with fibromyalgia. *Arthritis Rheum* 1986; 29: 1371-1377.

Gowans SE, De Hueck A, Voss S, et al. Effect of a randomized, controlled trial of exercise on mood and physical function in individuals with fibromyalgia. *Arthritis Rheum* 2001; 45: 519-529.

Gowans SE, De Hueck A, Voss S, Richardson M. A randomized, controlled trial of exercise and education for individuals with fibromyalgia. *Arthritis Care Res* 1999; 12: 120-128.

Häkkinen K, Paharinen Lakkinen A, Pakarinen P, Hannonen, Häkkinen A, Airaksinen O, Valkeinen H, Alen M. Effects of Strength Training on Muscle Strength, Cross-Sectional Area, Maximal Electromyographic Activity, and Serum Hormones in Premenopausal Women with Fibromyalgia. *J Rheumatol* 2002; 29: 1287-1295.

Iversen MD, Eaton HM, Daltroy LH. How rheumatologists and patients with rheumatoid arthritis discuss exercise and the influence of discussions on exercise prescriptions. *Arthritis Care Res* 2004; 51: 63-72.

Jentoft ES, Kvalvik AG, Mengshoel AM. Effects of pool-based and land-based aerobic exercise on women with fibromyalgia/chronic widespread muscle pain. *Arthritis Rheum* 2001; 45: 42-47.

Jones KD, Burckhardt CS, Clark SR, et al. A randomized controlled trial of muscle strengthening versus flexibility training in fibromyalgia. *J Rheumatol* 2002a, 29:1041-1048.

Jones KD, Clark SR. Individualizing the exercise prescription for persons with Fibromyalgia *Rheumatic Diseases Clinics of North America* 2002; 28 (2).

Keel PJ, Bodoky C, Gerhard U, Muller W. Comparison of integrated group therapy and group relaxation training for fibromyalgia. *Clin J Pain* 1998; 14: 232-238.

Krsnich-Shriwise S. Fibromyalgia syndrome: an overview in physical therapy. *Phys Ther.* 1997; 77: 68-75.

Lemstra M, Olszynski M and W. The effectiveness of multidisciplinary rehabilitation in the treatment of fibromyalgia. *Clin J Pain* 2005; 21(2): 166-174.

Mannerkopi K, Nyberg B, Ahlmen M, Ekdahl Ch. Pool exercise combined with an education program for patients with fibromyalgia syndrome. A prospective, randomized study. *J Rheumatol* 2000; 27: 2473-2481.

Mannerkopi K, Ahlmen M, Ekdahl C. Six- and 24-month follow-up of pool exercise therapy and education for patients with fibromyalgia. *Scand J Rheumatol* 2002; 31: 306-310.

Mannerkorpi K, Iversen MD. Physical exercise in fibromyalgia and related syndromes. *Best Pract Res Clin Rheumatol* 2003; 17: 629-647.

Mannerkorpi K. Exercise in fibromyalgia. *Curr Opin Rheumatol* 2005; 17 (2):190-194.

Martin L, Nutting A, Macintosh BR, Edworthy SM, Butterwick D, Cook J. An exercise program in the treatment of fibromyalgia. *J Rheumatol* 1996; 23: 1050-1053.

Martínez Lavín M. Management of Disautonomia in Fibromyalgia. *Rheum Dis Clin North Am* (2002); 28: 379-88.

McCain GA. A cost effective approach to the diagnosis and treatment of fibromyalgia. *Rheum Dis Clin North Am* 1996; 22: 323-49.

Meyer BB, Lemley KJ. Utilizing exercise to affect the symptomology of fibromyalgia: a pilot study. *Med Sci Sports Exerc* 2000; 32(10): 1691-1697.

Millea PJ, Holloway RL. Treating fibromyalgia. *American Family Physician*. 2000; 62(7):1575-82, 1587.

Ministerio de Sanidad y Consumo. Fibromialgia. Madrid: Ministerio de Sanidad y Consumo; 2003.

Monterde S, Salvat I, Montull S y Fernández Ballart J. Validación de la versión española del Fibromyalgia Impact Questionnaire. *Rev Esp Reumatol* 2004; 31 (9):507-513.

Moss-Morris R, Sharon C, Tobin R, Baldi, JC. A randomized controlled graded exercise trial for chronic fatigue syndrome: outcomes and mechanisms of change. *Journal of Health Psychology* 2005; 10(2): 245-259.

Moulin DE. Systemic drug treatment for chronic musculoskeletal pain. *Clin J Pain* 2001; 17: S86-93.

Muñoz M, Pérula LA, Ruiz R. Eficacia del método centrado en el paciente en sujetos con dolor crónico musculoesquelético generalizado no articular en Atención Primaria. *Accésit premio investigación en salud en Andalucía* 2001. Sevilla: Consejería de Salud; 2002.

Nicassio PM, Radojevic V, Weissman MH, Schuman C, Kim J, Schoenfeld-Smith K, Krall T. A Comparison of Behavioral and Educational Interventions for Fibromyalgia. *J Rheum* 1997; 24: 2000-7.

Nicassio PM, Shuman C, Kim J, Cordova A, Weissman MH. Psychosocial Factors Associated With Complementary Treatment Use in Fibromyalgia. *J Rheum* 1997; 24: 2008-13.

Offenbaecher M, Waltz M. Validation of a German Version of the Fibromyalgia Impact Questionnaire (FIQ-G). *Journal of Musculoskeletal Pain* Vol 6S; 1998: 90.

Pate R, Pratt NI, Blair SN, et al. Physical activity and public health: a recommendation from the Centers for Disease Control and Prevention and the American College of Sports Medicine. *JAMA* 1995; 273: 402-407.

Powell P, Bentall RP, Nye FJ, Edwards HT. Randomised controlled trial of patient education to encourage graded exercise in chronic fatigue syndrome. *BMJ* 2001; 322: 1-5.

Qué es la Fibromialgia. Sociedad Española de Reumatología [en línea] 2005; 24(1-6). URL disponible en: <http://www.ser.es/>

Ramsai C, Moreland J, Ho M, Joyce S, Walker S, Pullar T. An observer-blinded comparison of supervised and unsupervised aerobic exercise regimens in fibromyalgia. *Rheumatology* 2000; 39: 501-505.

Rao SG, Bennett RM. Pharmacological therapies in Fibromyalgia. *Best Practice&Research Clinical Rheumatology* 2003; 17 (4): 611-627.

Richards SCM, Scott DL. Prescribed exercise in people with fibromyalgia: parallel group randomised controlled trial. *BMJ* 2002; 325:185-188.

Rivera J, Gonzalez T. The Fibromyalgia Impact Questionnaire: a validated Spanish version to assess the health status in women with fibromyalgia. *Clin Exp Rheumatol.* 2004; 22(5): 554-60.

Rusell IJ, Kamin M, Bennett R, et al. Efficacy of tramadol in treatment of pain in fibromyalgia. *J Clin Rheumatol* 2000; 6: 250-7.

Schachter CL, Busch AJ, Peloso PM, et al. Effects of short versus long bouts of aerobic exercise in sedentary women with fibromyalgia: a randomized, controlled trial. *Phys Ther* 2003, 83: 340-358.

Selwyn CM, Richards, David L Scott. Prescribed exercise in people with fibromyalgia: parallel group randomised controlled trial. *BMJ* 2002; 325: 185.

Sociedad Española de Reumatología. Estudio EPISER. Madrid: Sociedad Española de Reumatología; 2000.

Sorensen J, Bengtsson A, Backman E, Henriksson KG, Bengtsson M. Pain analysis in patients with fibromyalgia. Effects of intravenous morphine, lidocaine, and ketamine. *Scand J Rheumatol* 1995; 24: 360-5.

Taggart HM., Arslanian CL, Bae S, Singh K. Effects of T'ai Chi exercise on fibromyalgia symptoms and health-related quality of life. *Orthop Nurs* 2003; 22 (5): 353-360.

The Copenhagen Declaration. Available from Bente Danneskiold-Sanmsoe, Department of Rheumatology, Frederiksberg Hospital, Ndr Fasanvej, DK-2000 Frederiksberg, Denmark.

Tiidus PM, Pierrynowski M, Dawson KA. Influence of moderate training gait and work capacity of fibromyalgia patients: a preliminary fields study. *Journal of Sports Science and Medicine* 2002; 1:122-127.

Tofferi JK, Jackson JL, O'Malley PG. Treatment of fibromyalgia with cyclobenzaprine: a meta-analysis. *Arthritis Rheum* 2004; 51: 9-13.

Valim V, Oliveira L, Suda A, et al. Aerobic fitness effects in fibromyalgia. *J Rheumatol* 2003; 30: 1060-1069.

Valverde M, Juan A, Rivas B, Carmona L. Estudio EPISER. Prevalencia e impacto de las enfermedades reumáticas en la población adulta española. Madrid: Editorial MSD y Sociedad Española de Reumatología; 2001.

Vlaeyen JWS, Teeken-Gruben NJG, Goossens MEJB, Rutten-van Mölken MPMH, Pelt RAGB, van Eek H, Heuts PHTG. Cognitive-Educational Treatment of Fibromyalgia: A randomized Clinical Trial. I. Clinical Effects. *J Rheum* 1996; 23: 1237-45.

White KP, Harth M. An analytical review of 24 controlled clinical trials for fibromyalgia syndrome (FMS). *Pain Vol* 1996; 64(2) 211-219.

WHO. International Statistical Classification of Diseases and Related Problems. ICD-10. Geneva: WHO, 1992.

Wigers SH, Stiles TC, Vogel PA. Effects of aerobic exercise versus stress management treatment in fibromyalgia. A 4.5 year prospective study. *Scand J Rheumatol* 1996; 25: 77-86.

Wolf F and The Vancouver Fibromyalgia Consensus Group. The Fibromyalgia Syndrome: A Consensus Report on Fibromyalgia and Disability. *J Rheum* 1996; 23: 534-39.

Wolf F, Anderson J, Harkness D, Bennet RM, Caro XJ, Goldenberg DL, Russel IJ, Yunus MB. Health Status and Disease Severity in Fibromyalgia. Results of a Six-Center Longitudinal Study. *Arthritis Rheum* 1997; 40: 1571-1579.

Wolf F, Anderson J, Harkness D, Bennet RM, Caro XJ, Goldenberg DL, Russell IJ, Yunus MB. Work and Disability Status of Persons with Fibromyalgia. *J Rheum* 1997; 24:1171-8.

Wolfe F, Anderson J, Harkness D, Bennet RM, Caso SJ, Goldenberg DL et al. A prospective, longitudinal, multicenter study of service utilization and costs in fibromyalgia. *Arthritis Rheum* 1997; 40(9): 1560-70.

Wolfe F, Smythe HA, Yunus MB, Bennett RM et al. The American College of Rheumatology 1990 Criteria for the Classification of Fibromyalgia. Report of the Multicenter Criteria Committee. *Arthritis Rheum* 1990; 33: 160-72.