



12

Bibliografía

1. Bone RC, Balk RA, Cerra FB et al. American College of Chest Physicians/Society of Critical Care Medicine. Consensus Conference: Definitions for sepsis and organ failure and guidelines for the use of innovative therapies in sepsis. *Crit Care Med* 1992; 20:864-874.
2. Rancel-Fausto MS, Pittet D, Costigan M et al. The natural history of the systemic Inflammatory Response Syndrome (SIRS). A prospective study. *JAMA* 1995; 273: 117-223.
3. Balk, RA. Severe sepsis and septic shock. Definitions, epidemiology, and clinical manifestations. *Crit Care Clin* 2000; 16:179-192.
4. Levy MM, Fink MP, Marshall JC, Abraham E, Angus D, Cook D et al. 2001 SCCM/ESICM/ACCP/ATS/SIS International Sepsis Definitions Conference. *Crit Care Med* 2003; 31:1250-1256.
5. Leon Gil C, Garcia-Castrillo Riesgo L, Moya Mir M et al. (Consensus document (SEMES-SEMICYUC). Recommendations for the initial and multidisciplinary diagnostic management of severe sepsis in the hospital Emergency Departments). *Med Intensiva* 2007; 31(7):375-387.
6. Dellinger RP, Levy MM, Carlet JM et al. Surviving Sepsis Campaign: international guidelines for management of severe sepsis and septic shock: 2008. *Crit Care Med* 2008; 36(1):296-327.
7. Carcillo JA, Fields AI, Task Force Comité Members, Clinical practice parameters for hemodynamic support of pediatric and neonatal patients in septic shock. *Crit Care Med* 2002; 30 (6): 1365-1378.
8. Calandra T, Cohen J; Internacional Sepsis Forum Definition of Infection in the ICU Consensus Conference. The international sepsis forum consensus conference on definitions of infection in the intensive care unit. *Crit Care Med* 2005; 33: 1538-1548.
9. Goldstein B, Giroir B, Randolph A

- and the Members of the International Consensus Conference on Pediatric Sepsis. International pediatric sepsis consensus conference: Definitions for sepsis and organ dysfunction in pediatrics. *Pediatr Crit Care Med* 2005; 6 (1):2-6.
10. Gebara BM, Beaumont W. Values for Systolic Blood Pressure. Letters to the Editor. (Goldstein 2005) *Pediatr Crit Care Med* 2005 Vol. 6, No. 4. 500.
11. Goldstein B, FCCM, Giroir B, Randolph A. Letters to the Editor *Pediatr Crit Care Med* 2005 Vol. 6, No. 4. 500.
12. NHLBI. National Heart, Lung, and Blood Institute. The Harriet Lane Handbook, 13th ed. Report of the second Task Force i Blod Pressure Control in Children. 1987. *Pediatrics*. 1987; 79:1-25.
13. Carrillo Álvarez A, et al. Reconocimiento del niño con riesgo de parada cardiorrespiratoria. *An Pediatr (Barc)*. 2006;65(2):147-53.
14. Pollack MM et al. Pediatric Risk of Mortality score (PRISM). *Crit Care Med* 1988;16:1110-6.
15. Leteurtre S, Martinot A, Duhamel A, et al. Validation of the paediatric logistic organ dysfunction (PELOD) score: prospective, observational, multicentre study. *Lancet* 2003; 362: 192-97.
16. Consejería de Salud, Junta de Andalucía. Estrategia para la Seguridad del Paciente.
17. LEY 41/2002, de 14 de noviembre, básica reguladora de la autonomía del paciente y de derechos y obligaciones en materia de información. BOE 427; 40126-40132.
18. Alonso Salas MT, de Carlos Vicente JC, Gil Antón J, Pinto Fuentes I, Quintilla Martínez JM, Sánchez Díaz JL. Documento de consenso SECIP-SEUP sobre manejo de sepsis grave y shock séptico en pediatría. http://www.secip.eu/05_infeccioso/ PROTOCOLOSECIPSEUPSEPSIS2009.pdf
19. Hazelzet JA. Diagnosing meningococemia as a cause of sepsis. *Pediatr Crit Care Med*. 2005 Vol. 6, No. 3 (Suppl.):S50-S54.
20. Proceso Asistencial Integrado Síndrome febril en el niño. Consejería de Salud. Junta de Andalucía. Sevilla. Consejería de Salud, 2003. 85 p.
21. Ochs HD. Steroids and immunizations. *Pediatr Infect Dis J*. 1994;13:760
22. Chaudhuri A, Martinez-Martin P, Kennedy PG, Andrew Seaton R, Portegies P, Bojar M, Steiner I, EFNS Task Force. EFNS guideline on the management of community-acquired bacterial meningitis: report of an EFNS Task Force on acute bacterial meningitis in older children and adults. *Eur J Neurol*. 2008 15):649-659.
23. Gervais A, Caffish M and Suter S. Management of children with fever without localizing signs of an infection. *Arch Pediatr*. 2001 8; 3: 324-330.
24. Bonadio WA, Hennes H, Smith D, et al. Reliability of observation variables in distinguishing infectious outcome of febrile young infants. *Pediatr Infect Dis J* 1993; 12:112-114.
25. RM, Hazday N, Pena M et al. Clinical antecedents to in-hospital-cardiopulmonary arrest. *Chest* 1990;98:1388-1392.
26. Franklin C, Mathew J. Developing strategies to prevent in-hospital cardiac arrest: analyzing responses of physicians and nurses in the hours before the event. *Crit Care Med* 1994;22:244-247.
27. ACEP: Clinical policy for children younger than three years presenting to the emergency department with fever. American College of Emergency Physicians - Medical Specialty Society. 2003 Oct.
28. Rivers E, Nguyen B, Havstad S, Ressler J, Muzzin A, Knoblich B, for the Early Goal-Directed Therapy Collaborative Group. Early goal-directed therapy in the treatment of severe sepsis and septic shock. *N Engl J Med*

2001 8;345:1368-1377.

29. Jones AE, Shapiro NI, Roshon M. Implementing early goal-directed therapy in the emergency setting: the challenges and experiences of translating research innovations into clinical reality in academic and community settings. *Acad Emerg Med.* 2007 Nov;14(11):1072-1078.

30. Lever A, Mackenzie I. Sepsis: definition, epidemiology, and diagnosis. *BMJ.* 2007 27;33, 7625:879-883.

31. Ley 5/2003, de 9 de octubre, de declaración de voluntad vital anticipada. BOE núm. 279:41231-41234.

32. Sinclair JF, Skeoch CH, D. Hallworth D. Prognosis of meningococcal septicaemia. *Lancet* 1987; 2(8549):38.

33. Bridgewater K. Caring for children with fever. RCN goodpractice guidance for nurses working with infant, children and young people. Published by the Royal College of Nursing, W1G ORN.2008 Royal. www.rcn.org.uk

34. <http://www.sepsisforum.org>

35. Johnson M, Maas, M, Moordhead S. (eds). *Nursing Outcomes Classification (NOC)*, Mosby, St. Louis 2000, 2ª ed.

36. Herrera Carranza M, Garcia Garmedia JL, Borrego Cobos J, et al. Plan Andaluz de Urgencias y Emergencias. Traslado de enfermos críticos. Protocolos de Transporte Secundario y Primario. Consejería de Salud, Junta de Andalucía, 2000.

37. Booy R, Habibi P, Nadel S, et al. Reduction in case fatality rate from meningococcal disease associated with improved healthcare delivery. *Arch Dis Child* 2001; 85: 386-390.

38. Bulechek GM, Butcher HK, McCloskey Dochterman J. *Clasificación de Intervenciones de Enfermería (NIC)*, Elsevier, 5ª ed. 2009.

39. Asociación Española de Nomenclatura, Taxonomía y Diagnósticos de Enfermería. AENTDE (Trad.). *Etiquetas y*

códigos de las intervenciones de Enfermería. 1995 Iowa Intervention Project.

40. Rivers E. The outcome of patients presenting to the emergency department with severe sepsis or septic shock. *Crit Care.* 2006;10(4):154-156.

41. Nee P, Andrews F, Rivers E. Critical care in the emergency department: introduction. *Emerg Med J.* 2006 Jul;23(7):560.

42. Rivers E. Implementation of an evidence-based "standard operating procedure" and outcome in septic shock: what a sepsis pilot must consider before taking flight with your next patient. *Crit Care Med.* 2006 Apr;34(4):1247.

43. Carcillo JA. Reducing the global burden of sepsis in infants and children: a clinical practice research agenda. *Pediatr Crit Care Med* 2005; 6 (3 Suppl): S157-S164.

44. Thomson APJ, Riordan FAI. Guidelines for the Management of Meningococcal Septicaemia. *Eur J Pediatr* 2002;161: 531-7.

45. Lin JC, Kuch BA, Felmet KA et al. Best practice outcome attained with implementation of ACCM/PALS guidelines for resuscitation of neonatal and pediatric septic shock; an UTSEIN Style análisis. *Crit Care Med.* 2003; 31 (12): A70.

46. Oliveira CF, Nogueira de Sa FR, Oliveira D, Gottschald A, Moura J, Shibata O, Troster E, Vaz F, MD, Carcillo JA. Time- and Fluid-Sensitive Resuscitation for Hemodynamic Support of Children in Septic Shock. Barriers to the Implementation of the American College of Critical Care Medicine/ Pediatric Advanced Life Support Guidelines in a Pediatric Intensive Care Unit in a Developing World. *Pediatric Emergency Care.* 2008; 24 (12): 810-8.

47. ORDEN de 8 de julio de 2009, por la que se dictan instrucciones a los Centros del Sistema Sanitario Público de Andalucía, en relación al procedimiento de Consentimiento Informado. BOJA nº152, 72-79.

48. http://www.juntadeandalucia.es/salud/sites/csalud/contenidos/Informacion_General/p_3_p_11_procedimiento_consentimiento_informado.
49. Brierley J, Carcillo JA, Choong K, Cornell T, DeCaen A, Deymann A et al. Clinical practice parameters for hemodynamic support of pediatric and neonatal septic shock: 2007 update from the American College of Critical Care Medicine*. *Crit Care Med* 2009; 37:666–688.
50. Muñoz Muñoz N, Alegre del Rey E, Díaz Navarro J, Morillo Verdugo R y Márquez Saavedra E (Editores). *Antiinfecciosos para uso sistémico*. En *Guía Farmacoterapéutica de Hospital (2008)*. (Pag.111-162) Sociedad de Farmacéuticos de Hospitales y Centros Socio Sanitarios. Sevilla.
51. Garnacho-Montero J, García-Garmendia JL, Barrero Almodovar A, Jiménez-Jiménez FJ, Pérez-Paredes C, Ortiz-Leyba C. Impact of adequate empirical antibiotic therapy on the outcome of patients admitted to the intensive care unit with sepsis. *Crit Care Med* 2003; 31: 2742-2751.
52. Garnacho-Montero J, Ortiz-Leyba C, Herrera-Melero I, Aldabo-Pallas T, Cayuela-Dominguez A, Marquez-Vacaro JA, Carbajal-Guerrero J, García-Garmendia JL. Mortality and morbidity attributable to inadequate empirical antimicrobial therapy in patients admitted to the ICU with sepsis: a matched cohort study. *J Antimicrob Chemother* 2008; 61:436-441.
53. Kumar A, Roberts D, Wood KE, Light B, Parrillo JE, Sharma S et al. Duration of hypotension before initiation of effective antimicrobial therapy is the critical determinant of survival in human septic shock. *Crit Care Med* 2006;34(6):1589-96.
54. Kollef MH. Inadequate antimicrobial treatment: an important determinant of outcome for hospitalized patients. *Clin Infect Dis* 2000; 31 (Suppl 4): S131-138.
55. Rodríguez Baño J, Navarro MD, Romero L, et al. Bacteremia due to extended-spectrum beta -lactamase-producing *Escherichia coli* in the CTX-M era: a new clinical challenge. *Clin Infect Dis*. 2006; 43:1407-1414.
56. Cisneros-Herreros JM, Cobo-Reinoso J, Pujol-Rojo M, Rodríguez-Baño J, Salavert-Lletí M. Guía para el diagnóstico y tratamiento del paciente con bacteriemia. *Guías de la Sociedad Española de Enfermedades Infecciosas y Microbiología Clínica (SEIMC)*. *Enferm Infecc Microbiol Clin* 2007 25:111-130.
57. Russell JA. Management of Sepsis. *N Eng J Med* 2006; 255:1699-713.
58. Anónimo. Treatment guidelines from the Medical Letter. Choice antibacterial drugs. Sepsis syndrome 2007; 5(57): 33-50.
59. Guía de Práctica Clínica. Diagnóstico y tratamiento de sepsis grave y choque séptico en el adulto. Centro Nacional de Excelencia Tecnológica en Salud. México. Visto en: <http://www.cenetec.salud.gob.mx/interior/gpc.html>. Acceso el 12/04/2010.
60. DynaMed Editorial Team. Aspiration pneumonia. Última actualización el 30/03/2010. Disponible en DynaMed: <http://www.ebscohost.com/dyna-med>. Acceso el 12/04/2010.
61. Bartlett JG. Aspiration pneumonia in adults. Última actualización el 16/01/2009. Disponible en: Uptodate: <http://www.uptodate.com> Acceso el 12/04/2010.
62. López-Herce Cid ¿Debe administrarse tratamiento antibiótico prehospitalario en la sospecha de sepsis meningocócica? *J. Evid Pediatr*. 2006; 2:65.
63. Schein M, Marshall J. Source Control for surgical infections. *World J Surg*, 2004; 28:638-645.
64. Bunt TJ. Non-directed relaparotomy for intraabdominal sepsis: a futile procedure. *Am Surg* 1986; 52:294-298
65. Laing FC: Ultrasonography of the acute abdomen. *Radiol Clin North Am*

1992;30:389.

66. Lichtenstein D, Axler O: Intensive use of general ultrasound in the intensive care unit. Prospective study of 150 consecutive patients. *Intensive Care Med* 1993;19; 353.

67. Loza Fernández de Bobadilla E, Planes Reig A, Rodríguez-Créixems M. 3ª.Hemocultivos 2003. En: Cercenado E, Cantón R, editores. Procedimientos en Microbiología Clínica. Recomendaciones de la Sociedad Española de Enfermedades Infecciosas y Microbiología Clínica. Disponible en: <http://www.seimc.org/protocolos/microbiologia>.

68. Bouza E, Sousa D, Rodríguez-Créixems M, García-Lechuz J, Muñoz P. Is the volume of blood cultured still a significant factor in the diagnosis of bloodstream infections? *J. Clin. Microbiol* 2007; 45: 2765-2769.

69. Lee A, Mirrett S, Reller LB, Weinstein MP. Detection of bloodstream infections in adults: How many blood cultures are needed? *J Clin Microbiol* 2007; 45: 3546-3548.

70. Falagas ME, Kazantzi MS, Bliziotis IA. 2008. Comparison of utility of blood culture from intravascular catheters and peripheral veins: a systematic review and decision analysis. *J Med Microbiol* 2008; 57:1-8.

71. Vincent JL, De Mendonca A, Cantraine F, et al. Use of the SOFA score to assess the incidence of organ dysfunction/failure in intensive care units: Results of a multicenter, prospective study. *Crit Care Med* 1998;26:1793-1800.

72. Moreno R, Vincent JL, Matos R, et al. The use of maximum SOFA score to quantify organ dysfunction/failure in intensive care. Results of a prospective, multicenter study. *Intensive Care Med* 1999; 25: 686-696.

73. Janssens U, Graf C, Graf J, et al. Evaluation of the SOFA score: a single-center experience of a medical intensive care unit in 303 consecutive patients with predominantly cardio-

vascular disorders. *Intensive Care Med* 2000; 26: 1037-1045.

74. Graciano AL, Balko JA, Rahn DS, Ahmad N, Giroir BP. The Pediatric Multiple Organ Dysfunction Score (P-MODS) : development and validation of an objective scale to measure the severity of multiple organ dysfunction in critically ill children. *Crit care Med*. 2005; 33(7):1484-91. <http://www.sfar.org/scores2/pmods2.html>

75. Iribarren-Diarasarria S, Latorre-García K, Muñoz-Martínez T, Poveda-Hernández Y, Dudagoitia-Otaolea JL, Martínez-Alutiz S et al. Limitación del esfuerzo terapéutico tras el ingreso en una Unidad de Medicina Intensiva. Análisis de factores asociados. *Med Intensiva*. 2007; 31: 68-72.

76. Cabré L, Solsona JF y grupo de trabajo de bioética de la SEMICYUC. Limitación del esfuerzo terapéutico en Medicina Intensiva. *Med Intensiva* 2002; 26: 304-311.

77. Esteban A, Gordo F, Solsona LF, Alía I, Caballero J, Bouza C et al. Withdrawing and withholding life support in the intensive care unit: a Spanish prospective multi-centre observational study. *Intensive Care Med* 2001; 27:1744-1749.

78. Martín Delgado MC, Cabré Pericas L, Javier Ruíz Moreno J, Blanch Torra L, Blanco Varela J, Castillo Suro F et al. Indicadores de Calidad Asistencial. Semicyuc, Madrid, 2005.

79. Dotor Gracia M, Fernández García E, Álvarez Benito M et al. Consejería de Salud. Guía de diseño y mejora continua de procesos asistenciales integrados . 2ª ed. Sevilla. Junta de Andalucía, 2009.

80. Brun-Buisson C: The epidemiology of the systemic inflammatory response. *Intensive Care Med* 2000, 26 Suppl 1:S64-S74.

81. Vincent JL, Rello J, Marshall J, Silva E, Anzueto A, Martin CD et al. International Study of the Prevalence and Outcomes of Infection in Intensive Care Units. *JAMA* 2009; 302(21): 2323-2329

82. Barriere SL, Lowry SF: An overview of mortality risk prediction in sepsis. *Crit Care Med* 1995; 23:376-393.
83. Grupo para el Estudio de la Infección en Urgencias. Estudio epidemiológico de las infecciones en el Área de urgencias. *Emergencias*. 2000;12:80-9.
84. Martínez Ortiz de Zárate M. Aspectos epidemiológicos de las infecciones en las Áreas de Urgencias. *Emergencias*. 2001;13: S44-S50.
85. Olivan Ballabriga AS, Agudo Pascual O, Agorreta Ruiz J, Pinillos Echevarría MA, Etxebarria Lus M, Moros Morgoñón MA, et al. La patología infecciosa en el Servicio de Urgencias de un hospital terciario. *Emergencias* 1998;10 Supl 1:22-5.
86. Carter C. Implementing the severe Sepsis care bundles outsider the ICU by outreach. *British Association of Critical Care Nurses, Nursing in Critical Care* 2007, 12(5): 225-230.
87. Hillman K, Chen J, Cretikos M, Bellomo R, Brown D, Doig G et al. Introduction of the medical emergency team (MET) system: a cluster-randomised controlled trial. *Lancet* 2005; 365: 2091-2097.
88. Lighthall GK, Markar S, Hsiung R. Abnormal vital signs are associated with an increased risk for critical events in US veteran inpatients. *Resuscitation* 2009; 80(11): 1264-1269.
89. Frost SA, Alexandrou E, Bogdanovski T. Unplanned admission to intensive care after emergency hospitalisation: Risk factors and development of a nomogram for individualising risk. *Resuscitation* 2009; 80(2): 224-230.
90. Iñigo J, Sendra JM, Díaz R, Bouza C, y Sarría-Santamera A. Epidemiología y costes de la sepsis grave en Madrid. *Estudio de altas hospitalarias. Med Intensiva*. 2006; 30:197-203.
91. Watson RS, Carcillo, JA. Scope and epidemiology of pediatric sepsis. *Crit CareMed* 2005; 6S: s3-s5.
92. Arnal LE, Stein F. Pediatric septic shock: Why has mortality decreased? The utility of goal-directed therapy. *Seminars in Pediatric Infectious Diseases*. 2003;14(2):165-172.
93. Montalbán G, López O. Parámetros Clínicos y soporte hemodinámico en el niño y neonato en shock séptico: actualización. *Revista Electrónica de Medicina Intensiva*. Artículo nº A100. Vol 9 nº 4, abril 2009.
94. Karapinar B, Lin JC, Carcillo JA. ACCM guidelines use, correct antibiotic therapy, and immune suppressant withdrawal are associated with improved survival in pediatric sepsis, severe sepsis, and septic shock. *Crit Care Med* 2004; 32:S573;A161.
95. Wells LC, JSmith JC, Weston VC, Collier J, Rutter N. The child with a non-blanching rash: how likely is meningococcal disease?. *Arch Dis Child* 2001;85:218-222.
96. Nielsen H E, Andersen E A, Andersen J, Böttiger B, Christiansen K M, Daugbjerg P, Larsen S O, Lind I, Nir M, Olofsson K. Diagnostic assessment of haemorrhagic rash and fever. *Arc Dis Chil*. 2001, 85: 160-165.
97. López-Herce J. Diagnóstico y tratamiento de la sepsis grave en el niño. *Revista Electrónica de Medicina Intensiva*. Artículo nº C30. Vol 4 nº 8, agosto 2004.
98. Ochs HD. Steroids and immunizations *Pediatr Infect Dis J* 1994;13:760.
99. Casado Flores, J. "Fiebre e hipotensión. Shock séptico". In: Casado J., Serrano A. "Urgencias y tratamiento del niño grave". Ediciones Ergon S.A. Madrid 2000. Pag. 90-94.
100. Larry K. Pickering, MD. *Red Book. Medical Trends*, SL. Barcelona. 2004.
101. Carcillo JA, Piva PP, Thomas NJ, Han YY, Lin JC, Orr RA. Shock and shock síndromes. Slonim AD, ° MM. *Pediatric Critical Care Medicine*. Ed. Lippincott William & Wilkins. Philadelphia 2006. p. 438-471.
102. Baraff, L. J.; Bass, J. W.; Fleis-

- her, G. R.; Klein, J. O.; McCracken, G. H.; Powell, K. R.; and Schriger, D. L.: Practice guideline for the management of infants and children 0 to 36 months of age with fever without source. Agency for Health Care Policy and Research. *Ann Emerg Med.* 1993; 22(7): 1198-210.
103. Ishimine P. The evolving approach to the young child who has fever and no obvious source. *Emerg Med Clin North Am.* 2007 Nov;25(4):1087-2115.
104. Evidence based clinical practice guideline for fever of uncertain source in children 2 to 36 months of age. Cincinnati Children's Hospital Medical Center - Hospital/ Medical Center. 2000 Feb (revised 2003 Oct 27; reviewed 2006 Aug).
105. Evidence based clinical practice guideline for fever of uncertain source in infants 60 days of age or less. Cincinnati Children's Hospital Medical Center - Hospital/Medical Center. 1998 May 15 (revised 2003 Jun; reviewed 2006 Sep).
106. Bonadio WA. Assessing patient clinical appearance in the evaluation of the febrile child. *Am J Emerg Med* 1995; 13: 321-326.
107. McCarthy PL, Sharpe MR, Spiesel SZ, et al. Observation scales to identify serious illness in febrile children. *Pediatrics* 1982; 70: 802-809 .
108. Simpson D, Reilly P. Paediatric Coma Scale. *Lancet* 1982; 2:450.
109. Reilly P, Simpson D et al- Assessing the conscious level in infants and young children: a paediatric version of the Glasgow Coma Scale. *Child's Nerv Syst* 1988;4:30-3.
110. Han JJ, Carcillo JA, Dragotta MA, Bills DM, Watson RS, Westerman ME, Orr RA. Early Reversal of Pediatric-Neonatal Septic Shock by Community Physicians Is Associated With Improved Outcome. *Pediatrics.* 2003; 112(4):793-9.
111. Khilnani P, Deopujari S, Carcillo J. Recent Advances in Sepsis and Septic Shock. *Indian J Pediatr* 2008; 75:821-830.
112. Casado J, Blanco A. Sepsis meningocócica y shock séptico: medidas eficaces, tratamientos controvertidos. *An. Esp. Ped.* 1999. 50 (4): 346-352.
113. GRADE working group. Grading quality of evidence and strength of recommendations. *BMJ,* 2004; 328:1490–1498.
114. Guyatt G, Gutterman D, Baumann MH, et al. Grading strength of recommendations and quality of evidence in clinical guidelines: report from an American College of Chest Physicians task force. *Chest,* 2006; 129:174–181.
115. Schünemann HJ, Jaeschke R, Cook DJ, et al.: ATS Documents Development and Implementation Committee. An Official ATS Statement: Grading the Quality of Evidence and Strength of Recommendations in ATS Guidelines and Recommendations. *Am J Respir Crit Care Med,* 2006; 174:605–614.
116. Buist MD, Jarmolowski E, Burton PR, Bernard SA, Waxman. BP, Anderson J. Recognising clinical instability in hospital patients before cardiac arrest or unplanned admission to intensive care. A pilot study in a tertiary care hospital. *Med J Aust* 1999; 171:22-25.
117. Buist MD, Moore GE, Bernard SA, Waxman BP, Anderson JN, Nguyen TV. Cardiac arrests in hospital: preliminary study of incidence of and mortality from unexpected. *BMJ* 2002;324(7334);387-390.
118. Mackenzie I, Lever A. Management of sepsis. *BMJ.* 2007 Nov 3;335(7626):929-32.
119. Hollenberg SM. Vasopressor support in septic shock. *Lancet* 2007; 370(9602):1827-1828.
120. Keddissi JI, Bright BC, Dernaika TA. Use of pressors in the management of septic shock. *Lancet* 2007; 370:676-684.

121. Dries DJ. Vasoactive drug support in septic shock. *Shock*. 2006 Nov;26(5):529-30.
122. Oberbeck R. Catecholamines: physiological immunomodulators during health and illness. *Curr Med Chem*. 2006;13(17):1979-89.
123. Hou LC, Li SZ, Xiong LZ, et al. Effect of dopamine and metaraminol on the renal function of patients with septic shock. *Chin Med J (Engl)*. 2007 Apr 20;120(8):680-3.
124. Bellomo R, Chapman M, Finfer S, Hickling K, Myburgh J; Australian and New Zealand Intensive Care Society (ANZICS) Clinical Trials Group. Low-dose dopamine in patients with early renal dysfunction: a placebo controlled randomised trial. Australian and New Zealand Intensive Care Society (ANZICS) Clinical Trials Group. *Lancet* 2000; 356:2139–2143.
125. Kellum J, Decker J. Use of dopamine in acute renal failure: a meta-analysis. *Crit Care Med* 2001, 29:1526–1531.
126. Regnier B, Rapin M, Gory G, Lemaire, F, Teisseire B, Harari, A. Haemodynamic effects of dopamine in septic shock. *Intensive Care Med* 1977; 3:47–53.
127. Patel BM, Chittock DR, Russell JA, Walley KR. Beneficial effects of short-term vasopressin infusion during severe septic shock. *Anesthesiology* 2002, 96:576–582.
128. Dunser MW, Mayr AJ, Ulmer H, Knotzer H, Sumann G, Pajk W. Arginine vasopressin in advanced vasodilatory shock: a prospective, randomized, controlled study. *Circulation* 2002; 107:2313–2319.
129. Holmes CL, Patel BM, Russell JA, Walley KR. Physiology of vasopressin relevant to management of septic shock. *Chest* 2001, 120:989–1002.
130. Malay MB, Ashton RC, Landry DW, Townsend RN (1999) Low-dose vasopressin in the treatment of vasodilatory septic shock. *J Trauma* 1999, 47:699–705.
131. Holmes CL, Walley KR, Chittock DR et al. The effects of vasopressin on hemodynamics and renal function in severe septic shock: a case series. *Intensive Care Med* 2001, 27:1416–1421.
132. Lauzier F, Levy B, Lamarre P, Lesur O. Vasopressin or norepinephrine in early hyperdynamic septic shock: a randomized clinical trial. *Intensive Care Med* 2006, 32:1782–1789
133. O'Brien A, Calpp L, Singer M. Terlipressin for norepinephrine-resistant septic shock. *Lancet* 2002, 359:1209–1210.
134. Djillali A, Vigno P, Renault A and for the CATS STUDY Group (2007) Norepinephrine plus donutamine versus epinephrine alone for management of septic shock: A randomized trial. *Lancet* 2007,370:676–684.
135. Gattinoni L, Brazzi L, Pelosi P et al. A trial of goal-oriented hemodynamic therapy in critically ill patients. *N Engl J Med* 1995, 333:1025–1032
136. Hayes MA, Timmins AC, Yau EH et al. Elevation of systemic oxygen delivery in the treatment of critically ill patients. *N Engl J Med* 330:1717–1722.
137. Olaecha PM, Alvarez-Lerma F, Sanchez M. Evaluación estado de pacientes con infecciones graves. Criterios de ingreso en unidades de intensivo. *Revista: Enfermedades infecciones microbiología clinica* 2009. vol 27 pag 342 a 352.
138. Cooper DJ, Walley KR, Wiggs BR, et al. Bicarbonate does not improve hemodynamics in critically ill patients who have lactic acidosis: a prospective, controlled clinical study. *Ann Intern Med* 1990, 112:492–498.
139. Jiménez MF, Marshall JC. Source control in the management of sepsis. *Intensive Care Med* 2000; 27:S49-S62.
140. Schoeffel U. Timing of intervention. In Schein M, Marshall JC, (ed) *Source Control*. Heidelberg. Springer-Verlag. 2002.

141. Majeski JA, Alexander JW. Early diagnosis, nutritional support, and immediate extensive debridement improve survival in necrotizing fasciitis. *Am J Surg* 1983; 145(6):784-787.
142. Miller JD. The importance of early diagnosis and surgical treatment of necrotizing fasciitis. *Surg Gynecol Obstet*. 1983 ;157(3):197-200.
143. Sudarsky LA, Laschinger JC, Coppa GF, Spencer FC. Improved results from a standardized approach in treating patients with necrotizing fasciitis. *Ann Surg*. 1987; 206(5):661-665.
144. Moss RL, Musemeche CA, Kosloke AM. Necrotizing fasciitis in children: prompt recognition and aggressive therapy improve survival. *J Pediatr Surg*, 1996; 31:1142-1146.
145. Gang GI, Moulton JF, Solomkin JS. Drainage. In Schein M, Marshall JC, (ed) *Source Control*. Heidelberg. Springer-Verlag. 2002.
146. Poret HA, Fabian TC, Croce MA, et al. Analysis of septic morbidity following gunshot wounds to the colon: the missile is and adjuvant for abscess. *J. Trauma Injury Infect. Crit Care* 1999; 31:1088-1095.
147. Bohnen J. Debridement and "peritoneal toilet". In Schein M, Marshall JC, (ed) *Source Control*. Heidelberg. Springer-Verlag. 2002.
148. Evans A, Winslow BH. Oxygen saturation and hemodynamic response in critically ill mechanically ventilated adults during intrahospital transport. *Am J Crit Care*, 1995; 4:106-111.
149. Gerzof SG, Johnson WC, Robbins AH, Nabseth DC. Expanded criteria for percutaneous abscess drainage. *Arch Surg* 1985; 120(2):227-232.
150. Bufalari A, Giustozzi G, Moggi L. Postoperative intraabdominal abscesses: Percutaneous versus surgical treatment. *Acta Chir Bel*, 1996; 96:197-200.
151. Dunn HL. The biological rationale of source control. In Schein M, Marshall JC, (ed) *Source Control*. Heidelberg. Springer-Verlag. 2002.
152. Barie PS, Eachempati SR. Failed source control. In Schein M, Marshall JC, (ed) *Source Control*. Heidelberg. Springer-Verlag. 2002.
153. Mier J, Leon FL, Castillo A, Robledo F, Blanco R. Early versus Late necrosectomy in severe necrotizing pancreatitis. *Am J Surg* 1997; 173(2):71-71.
154. Knochel JQ, Koehler PR, Lee TG, Welch DM. Diagnosis of abdominal abscesses with computed tomography, ultrasound and 111In leukocyte scans. *Radiology* 1980; 137(2):425-432.
155. Schein M, Hirshberg A, Hashmonai M. Current surgical management of severe intrabdominal infection. *Surgery* 1992;112(3):489-496.
156. Hallerback B, Andersson C, Englund N et al. A prospective randomized study of continuous peritoneal lavage postoperatively in the treatment of purulent peritonitis. *Surg Gynecol Obstet* 1986; 163(5):433-436.
157. Cook DJ, Randolph A, Kenerman P, Cupido C, King D, Soukup C. Central venous catheter replacement strategies: a systematic review of the literature. *Cri Car Med* 1997; 25:1417-1424.
158. Bregenzer T, Conen D, Sakmann P, Widmer AF. Is routine replacement of peripheral intravenous catheters necessary? *Arch Inter Med* 1998; 158(2):151-156.
159. O'Grady NP, Alexander M, Dellinger EP, Gerberding JL, Heard SO, Maki DG et al. Guidelines for the prevention of intravascular catheter-related infections. *Centers for Disease Control and Prevention*, 2002; *MMWR* 51(RR-10):1-29.
160. O'Grady NP, Alexander M, Dellinger EP, Gerberding JL, Heard SO, Maki DG et al. Healthcare Infection Control Practices Advisory Committee. Guidelines for the prevention of intravascular catheter-related infections. *Clin Infect Dis*, 2002; 35:1281-1307
161. The Acute Respiratory Distress Syndrome Network. Venti-lation with

lower tidal volumes as compared with traditional tidal volumes for acute lung injury and the acute respiratory distress syndrome. *N Engl J Med* 2000, 342:1301–1308.

162. Amato MB, Barbas CS, Medeiros DM, Schettino Gde P, Lorenzi Filho G, Kairalla RA et al. Effect of a protective-ventilation strategy on mortality in the acute respiratory distress syndrome. *N Engl J Med* 1998; 338:347–354.

163. Brower RG, Shanholtz CB, Fessler HE, Shade DM, White PJ, Wiener CM et al. Prospective, randomized, controlled clinical trial comparing traditional versus reduced tidal volume ventilation in acute respiratory distress syndrome patients. *Crit Care Med* 1999; 27:1492–1498.

164. Stewart TE, Meade MO, Cook DJ, Granton JT, Hodder RV, Lapinsky SE et al. Evaluation of a ventilation strategy to prevent barotrauma in patients at high risk for acute respiratory distress syndrome. *N Engl J Med* 1998; 338:355–361.

165. Eichacker PQ, Gerstenberger EP, Banks SM, Cui X, Natanson C. Meta-analysis of acute lung injury and acute respiratory distress syndrome trials testing low tidal volumes. *Am J Respir Crit Care Med* 2002, 166:1510–1514.

166. Tobin MJ. Culmination of an era in research on the acute respiratory distress syndrome. *N Engl J Med* 2000, 342:1360–1361.

167. Marini JJ, Gattinoni L. Ventilatory management of acute respiratory distress syndrome: A consensus of two. *Crit Care Med* 2004, 32:250–255.

168. Hager DN, Krishnan JA, Hayden DL, Brower RG; ARDS Clinical Trials Network. Tidal volume reduction in patients with acute lung injury when plateau pressures are not high. *Am J Respir Crit Care Med* 2005; 172:1241–1245.

169. Ferguson ND, Frutos-Vivar F, Esteban A, and Study Group. Airway pressures, tidal volumes, and mortality in patients with acute respiratory distress

syndrome *Crit Care Med* 2005, 33:21–30.

170. Hickling KG, Henderson S, Jackson R. Low mortality rate in adult respiratory distress syndrome using low-volume, pressure-limited ventilation with permissive hypercapnia: a prospective study. *Crit Care Med* 1994, 22:1568–1578.

171. Bidani A, Tzouanakis AE, Cardenas VJ Jr, Zwischenberger JB. Permissive hypercapnia in acute respiratory failure. *JAMA* 1994, 272:957–962.

172. Kallet RH, Jasmer RM, Luce JM, Lin LH, Marks JD. The treatment of acidosis in acute lung injury with tris-hydroxymethyl aminomethane (THAM). *Am J Respir Crit Care Med* 2000, 161(4 Pt 1):1149–1153.

173. Weber T, Tschernich H, Sitzwohl C, Ullrich R, Germann P, Zimpfer M et al. Tromethamine buffer modifies the depressant effect of permissive hypercapnia on myocardial contractility in patients with acute respiratory distress syndrome. *Am J Respir Crit Care Med* 2000, 162:1361–1365.

174. Marini JJ, Ravenscraft SA. Mean airway pressure: physiologic determinants and clinical importance—Part 1: Physiologic determinants and measurements. *Crit Care Med* 1992, 20:1461–1472.

175. Gattinoni L, Marcolin R, Caspani ML, Fumagalli R, Mascheroni D, Pesenti A et al. Constant mean airway pressure with different patterns of positive pressure breathing during the adult respiratory distress syndrome. *Bull Eur Physiopathol Respir* 1985; 21:275–279.

176. Pesenti A, Marcolin R, Prato P, Borelli M, Riboni A, Gattinoni L. Mean airway pressure vs. positive end-expiratory pressure during mechanical ventilation. *Crit Care Med* 1985, 13:34–37.

177. The National Heart, Lung and Blood Institute ARDS Clinical Trials Network. Higher versus lower positive end-expiratory pressures in patients with the acute respiratory distress syndrome. *N Engl J Med* 2004, 351:327–336.

178. Villar J, Kacmarek RM, Pérez-Méndez L, Aguirre-Jaime A for the ARIES Network. A high PEEP-low tidal volume ventilatory strategy improves outcome in persistent ARDS. A randomized controlled trial. *Crit Care Med* 2006 34:1311–1318.
179. Amato MB, Barbas CS, Medeiros DM, Schettino Gde P, Lorenzi Filho G, Kairalla RA et al. Beneficial effects of the “open lung approach” with low distending pressures in acute respiratory distress syndrome. A prospective randomized study on mechanical ventilation. *Am J Respir Crit Care Med* 1995; 152:1835–1846.
180. Gattinoni L, Caironi P, Cressoni M, Chiumello D, Ranieri VM, Quintel M et al. Lung recruitment in patients with acute respiratory distress syndrome. *N Engl J Med* 2006; 354:1775–1786
181. Stocker R, Neff T, Stein S, Ecknauer E, Trentz O, Russi E et al. Prone positioning and low-volume pressure-limited ventilation improve survival in patients with severe ARDS. *Chest* 1997, 111:1008–1017.
182. Lamm WJ, Graham MM, Albert RK. Mechanism by which prone position improves oxygenation in acute lung injury. *Am J Respir Crit Care Med* 1994 150:184–193.
183. Jolliet P, Bulpa P, Chevrolet JC. Effects of the prone position on gas exchange and hemodynamics in severe acute respiratory distress syndrome. *Crit Care Med* 1998 26:1977–1985.
184. Gattinoni L, Tognoni G, Pesenti A for Prone-Supine Study Group. Effect of prone positioning on the survival of patients with acute respiratory failure. *N Engl J Med* 2001 345:568–573.
185. Guerin C, Gaillard S, Lemasson S, Ayzac L, Girard R, Beuret P, et al. Effects of systematic prone positioning in hypoxemic acute respiratory failure: a randomized controlled trial. *JAMA* 2004; 292:2379–2387.
186. Mancebo J, Fernández R, Blanch L, et al. A multicenter trial of prolonged prone ventilation in severe acute respiratory distress syndrome. *Am J Respir Crit Care Med* 2006;173:1233–1239.
187. Drakulovic MB, Torres A, Bauer TT, Nicolas JM, Nogue S, Ferrer M. Supine body position as a risk factor for nosocomial pneumonia in mechanically ventilated patients: a randomized trial. *Lancet* 1999; 354:1851–1858.
188. van Nieuwenhoven CA, Vandembroucke-Grauls C, van Tiel FH, van Schijndel, RJ, van der Tweel I, Ramsay G, et al. Feasibility and effects of the semirecumbent position to prevent ventilator-associated pneumonia: a randomized study. *Crit Care Med* 2006; 34:396–402.
189. Antonelli M, Conti G, Rocco M, Bufi M, De Blasi RA, Vivino G, et al. Mechanical ventilation in patients with acute respiratory failure. *N Engl J Med* 1998; 339:429–435.
190. Ferrer M, Esquinas A, Leon M, Gonzalez G, Alarcon A, Torres A et al. Noninvasive ventilation in severe hypoxemic respiratory failure: a randomized clinical trial. *Am J Respir Crit Care Med* 2003; 168:1438–1444.
191. Ely EW, Baker AM, Dunagan DP, Burke HL, Smith AC, Kelly PT et al. Effect on the duration of mechanical ventilation of identifying patients capable of breathing spontaneously. *N Engl J Med* 1996; 335:1864–1869.
192. Esteban A, Alía I, Tobin MJ, Gil A, Gordo F, Vallverdú I et al. Effect of spontaneous breathing trial duration on outcome of attempts to discontinue mechanical ventilation. *Am J Respir Crit Care Med* 1999; 159:512–518.
193. Esteban A, Alía I, Gordo F, Fernandez R, Solsona JF, Vallverdu I et al. Extubation outcome after spontaneous breathing trials with T-tube or pressure support ventilation. *Am J Respir Crit Care Med* 1997; 156:459–465.
194. Brochard L, Rauss A, Benito S, Conti G, Mancebo J, Rekić N et al. Comparison of three methods of gradual withdrawal from ventilatory support during weaning from mechanical ventilation. *Am J Respir Crit Care Med*

1994;150:896–903.

195. Cuthbertson BH, Sprung CL, Annane D, Chevret S, Garfield M, Goodman S et al. The effects of etomidate on adrenal responsiveness and mortality in patients with septic shock. *Intensive Care Med* 2009; 35:1868-1876.

196. Bernard GR, Vincent JL, Laterre PF, LaRosa SP, Dhainaut JF, Lopez-Rodriguez A et al. Recombinant human protein C Worldwide Evaluation in Severe Sepsis (PROWESS) study group. Efficacy and safety of recombinant human activated protein c for severe sepsis. *N Engl J Med* 2001; 344:699–709.

197. Abraham E, Laterre PF, Garg R, Levy H, Talwar D, Trzaskoma BL et al. Administration of Drotrecogin Alfa (Activated) in Early Stage Severe Sepsis (ADDRESS) Study Group. Drotrecogin alfa (activated) for adults with severe sepsis and a low risk of death. *N Engl J Med* 2005; 353:1332–1341.

198. Vincent JL, Bernard GR, Beale R, Doig C, Putensen C, Dhainaut JF et al. Drotrecogin alfa (activated) treatment in severe sepsis from the global open-label trial ENHANCE: Further evidence for survival and safety and implications for early treatment. *Crit Care Med* 2005; 33:2266–2277.

199. Oxman AD, Guyatt GH. A consumer's guide to subgroup analyses. *Ann Intern Med* 1992;116:78–84

200. Ely EW, Laterre PF, Angus DC, Helterbrand JD, Levy H, Dhainaut JF et al. Drotrecogin alfa (activated) administration across clinically important subgroups of patients with severe sepsis. *Crit Care Med* 2003;31:12–19.

201. Kanji S, Perreault MM, Chant C, Williamson D, Burry L. Evaluating the use of Drotrecogin alfa activated in adult severe sepsis: a Canadian multicenter observational study. *Intensive Care Med* 2007;33:517–523.

202. Bertolini G, Rossi C, Anghileri A, Livigni S, Addis A, Poole D. Use of Drotrecogin alfa (activated) in Italian intensive care units: The results of a nationwide survey. *Intensive Care Med*

2007;33:426–434.

203. European Medicines Agency. Evaluation of Medicines for Human Use. www.emea.europa.eu/pdfs/human/press/pr/8509607en.pdf. Accessed October 22, 2007.

204. Vieillard-Baron A, Caille V, Charron C et al. Reversal of refractory septic shock with drotrecogin alpha (activated). *Intensive Care Med* 2009; 39:1204-1209.

205. Knaus Wa, Draper EA, Wagner DP, Zimmerman JE. APACHE II: A severity of disease classification system. *Crit Care Med* 1985; 13: 818-829.

206. Finfer S. Corticosteroids in septic shock. *N Engl J Med*. 2008 Jan 10;358(2):188-90.

207. Sprung CL, Annane D, Keh D, Moreno R, Singer M, Freivogel K; COR-TICUS Study Group. Hydrocortisone therapy for patients with septic shock. *N Engl J Med* 2008 Jan 10; 358(2):111-124.

208. Irazuzta J, Sullivan KJ, Garcia PC, et al. Pharmacologic support of infants and children un septic shock. *J Pediatr* 2007;83(2 suppl): S36-45.

209. Parker MM, Hazelzet JA, Carcillo JA. Pediatric considerations. *Crit Care Med* 2004; 32 (11) (suppl): S591-S594.

210. Kanter RK, Zimmerman JJ, Strauss RH. Pediatric emergency intravenous access: Evaluation of a protocol. *Am J Dis Child* 1986; 140: 132-134.

211. Sempere AP, Berenguer-Ruiz L, Lezciano-Rodas M, Mira-Berenguer F, Waez M. Punción lumbar: indicaciones, contraindicaciones, complicaciones y técnica de realización. *Rev Neurol* 2007; 45: 433-436.

212. Kneen R, Solomon T, Appleton R. The role of lumbar puncture in suspected CNS infection a disappearing skill? *Arch Dis Child* 2002;87:181-183.

213. Riordan FAI et al. Guidelines for the Management of Meningococcal Septicaemia. *Eur J Pediatr* 2002; 161: 531-7. Review Date - June 2007

214. AHA Part 12: Pediatric Advanced Life Support. *Circulation* 2005; IV-167-IV-187. DOI: 10.1161/CIRCULATIONAHA.105.166573. http://circ.ahajournals.org/cgi/content/full/112/24_suppl/IV-167.
215. The International Liaison Committee on Resuscitation (ILCOR) Consensus on Science With Treatment Recommendations for Pediatric and Neonatal Patients: Pediatric Basic and Advanced Life Support. *Pediatrics* 2006; 117:e955-e977; DOI: 10.1542/peds.2006-0206 <http://www.pediatrics.org/cgi/content/full/117/5/e955>.
216. Guía del Servicio de Microbiología. Hospital Universitario Virgen de las Nieves. Granada. Mayo 2008.
217. Horstkotte D, Follath F, Gutschik E, Legyel M, Oto A, Pavia A et al. Guidelines on Prevention, Diagnosis and Treatment of Infective Endocarditis Executive Summary The Task Force on Infective Endocarditis of the European Society of Cardiology. *European Heart Journal* 2004; 25:267-276.
218. Connell TG, Rele M, Coeley D, Buttery JP, Curtis N. How reliable is a negative blood culture result) Volume of blood submitted for culture in routine practice in a children's hospital. *Pediatrics* 2007; 119:924-935.
219. Mermel LA, Allon M, Bouza E et al. Clinical Practice Guidelines for the Diagnosis and Management of Intravascular Catheter-Related Infection: 2009 Update by the Infectious Diseases Society of America. *IDSA Guidelines for Intravascular Catheter-Related Infection. Clinical Infectious Diseases* 2009; 49:1-45.
220. Ceneviva G, Paschall JA, Maffei F, Carcillo A. Hemodynamic support in fluid refractory pediatric septic shock. *Pediatrics* 1998; 102; 19. <http://www.pediatrics.org/cgi/content/full/102/2/e19>.
221. Boluyt N, Bollen CW, Bos AP, Kok JH, Offringa M. Fluid resuscitation in neonatal and pediatric hypovolemic shock: a Dutch Pediatric Society evidence-based clinical practice guideline. *Intensive Care Med.* 2006 Jul;32(7):958-961.
222. Stoner MJ, Goodman DG, Cohen DM, Fernandez SA, Hall MW. Rapid fluid resuscitation in pediatrics: testing the American College of Critical Care Medicine guideline. *Ann Emerg Med.* 2007 Nov;50(5):601-607.
223. Tabbutt S. Heart failure in pediatric septic shock: Utilizing inotropic support. *Crit Care Med* 2001; 29 (10) S231-S236.
224. Jackson WL Jr. Should we use etomidate as an induction agent for endotracheal intubation in patients with septic shock?: a critical appraisal. *Chest.* 2005;127:1031-1038.
225. Feltes T, Pignatelli R, Kleinert S, Mariscalco M. Quantitated left ventricular systolic mechanics in children with septic shock utilizing noninvasive wall stress analysis. *Crit Care Med.* 1994; 22: 1647-1658.
226. Michard F, Ayala S, Zarka V, Bahloul M, Richard C, Telboul JL. Global end diastolic volumen as an indicator of cardiac preload in patients with septic shock. *Chest.* 2003; 124:1900-1908.
227. Pollard AJ, Britto J, Nadel S. Emergency management of meningococcal disease. *Arch Dis Child* 1999; 80: 290-296.
228. Lacroix J, Hébert PC, Hutchison JS, Hume HA, Tucci M, Ducruet T, Gauvin F, Collet JP, Toledano BJ, Robillard P, Joffe A, Biarent D, Meert K, Peters MJ; TRIPICU Investigators; Canadian Critical Care Trials Group; Pediatric Acute Lung Injury and Sepsis Investigators Network. Transfusion strategies for patients in pediatric intensive care units. *N Engl J Med.*; 356(16):1609-19.
229. Annane D, Bellissant E, Bollaert PE, Briegel J, Keh D, Kupfer Y. Corticosteroides para el tratamiento de la sepsis grave y el shock séptico (Revisión Cochrane traducida). En: La Biblioteca Cochrane Plus, 2007 Número 2. Oxford: Update Software Ltd

230. Sarthi M, Lodha R, Vivekanandhan S, Arora NK. Adrenal status in children with septic shock using low-dose stimulation test. *Pediatr Crit Care Med*. 2007 Jan;8(1):23-8.
231. Guidet B, Mosqueda GJ, Priol G, Aegerter P. The COASST study: cost-effectiveness of albumin in severe sepsis and septic shock. *J Crit Care*. 2007 Sep; 22(3):197-203.
232. Maitland K, Pamba A, English M, Peshu N, Marsh K, Newton C. Randomized trial of volume expansion with albumin or saline in children with severe malaria: Preliminary evidence of albumin benefit. *Clin Infect Dis* 2005; 40:538-545.
233. Jacob M, Chappell D, Conzen P, Wilkes M M, Becker B F, Rehm M. Small-volume resuscitation with hyperoncotic albumin: a systematic review of randomized clinical trials. *Critical Care* 2008; 12:R34.
234. Demirdjam G. Albumina su uso basado en la evidencia científica, *Boletín CIME* 2003, VI (30)
235. de Oliveira CF, de Oliveira DS, Gottschald AF, Moura JD, Costa GA, Ventura AC et al. ACCM/PALS haemodynamic support guidelines for paediatric septic shock: An outcomes comparison with and without monitoring central venous oxygen saturation. *Intensive Care Med* 2008; 34: 1065-1075.
236. Lindsay CA, Barton P, Lawless S, Kitchen L, Zorka A, Garcia J et al. Pharmacokinetics and pharmacodynamic of milrinone lactate in pediatric patients with septic shock. *J Pediatr* 1998; 132:329-334.
237. Leit J, Jacqueline C, Orsonneau JL. The effects of milrinone on hemodynamics in experimental septic shock model. *Pediatric Crit Care Med* 2005; 6: 195-199.
238. De Luca L, Colucci WS, Nieminen MS, Massie BM, Gheorghide M Evidence-based use of Levosimendan in different clinical settings. *European Heart Journal* (2006) 27, 1908–1920.
239. Matok I, Leibovitch L, Vardi A, Adam M, Rubinshtein M, Barzilay Z, Paret G. Terlipressin as rescue therapy for intractable hypotension during neonatal septic shock. *Pediatr Crit Care Med* 2004 Mar;5(2):116-8.
240. Delmas A, Leone M, Rousseau S, Albanèse J, Martin C. Clinical review: Vasopressin and terlipressin in septic shock patients. *Crit Care*. 2005 Apr; 9(2): 212-22.
241. Rodríguez-Núñez A, López-Herce J, Gil-Antón J, Hernández A, Corsino and the RETSPED Working Group of the Spanish Society of Pediatric Intensive Care. Rescue treatment with terlipressin in children with refractory septic shock: a clinical study. *Critical Care* 2006, 10 (1): :R20 (doi:10.1186/cc3984).
242. Papoff P, Mancuso M, Barbara CS, Moretti C. The role of terlipressin in pediatric septic shock: a review of the literature and personal experience. *Int J Immunopathol Pharmacol*. 2007 Apr-Jun;20(2):213-21.
243. Rodriguez-Nuñez A., Fernandez-Sanmartin M, Martinon-Torres F, Gonzalez-Alonso N, Martinon-Sanchez JM. Terlipressin for catecholamine-resistant septic shock in children. *Intensive Care Med* 2004 Mar; 30(3):477-80.
244. Beale RJ, Hollemberg SM, Vincent JL, Parrillo JE. Vasopressor and inotropic support in septic shock: an evidence-based review. *Crit Care Med*. 2004; 32: S455-65.
245. Cooper BE. Review and Update on Inotropes and vasopressors. *AACN Advanced Critical Care*. 2008. 19 (1):5-10.
246. Fortenberry JD, Paden ML. Extracorporeal therapies in the treatment of sepsis: experience and promise. *Semin Pediatr Infect Dis*. 2006 Apr;17(2):72-9.
247. Leone M, Martin C. Role of terlipressin in the treatment of infants and neonates with catecholamine-resistant septic shock. *Best Pract Res Clin Anaesthesiol* 2008; Jun; 22(2):323-33.1

248. Meyer S, Gortner L, McGuire W, Baghai A, Gottschling S. Vasopressin in catecholamine-refractory shock in children. *Anaesthesia* 2008 Mar; 63(3):228-34.
249. Yildizdas D, Yapicioglu H, Celik U, Sertdemir Y, Alhan E. Terlipressin as a rescue therapy for catecholamine-resistant septic shock in children. *Intensive Care Med*. 2008 Mar; 33(3):511-7.
250. Fineman LD, LaBrecque MA, Shih MC, Curley MA. Prone positioning can be safely performed in critically ill infants and children. *Pediatr Crit Care Med*. 2006 Sep; 7(5):413-22.
251. Dahlem P, van Aalderen WM, Bos AP. Pediatric acute lung injury. *Paediatr Respir Rev*. 2007; 8(4):348-62.
252. Binld L, Nicolai T. Management of septic shock and acquired respiratory distress syndrome in pediatric cancer patients. *Klin Padriatr*. 2005, 217 Suppl 1:S130.
253. Branco RG, Garcia PC, Piva JP, Casartelli CH, Seibel V, Tasker RC. Glucose level and risk of mortality in pediatric septic shock. *Pediatr Crit Care Med*. 2005;6:470-2.
254. Srinivasan V, Spinella PC, Drott HR, Roth CL, Helfaer MA, Nadkarni V. Association of timing, duration, and intensity of hyperglycemia with intensive care unit mortality in critically ill children. *Pediatr Crit Care Med*. 2004; 5:329-36.
255. Van Waardenburg DA, Jansen TC, Vos GD, Buurman WA. Hyperglycemia in Children with Meningococcal Sepsis and Septic Shock: The Relation between Plasma Levels of Insulin and Inflammatory Mediators. *The Journal of Clinical Endocrinology & Metabolism*. 2006; 91(10): :3916–3921.
256. Kong MY, Alten J, Tofil N. Is hyperglycemia really harmful? A critical appraisal of "Persistent hyperglycemia in critically ill children" by Faustino and Apkon (*J Pediatr* 2005; 146:30-34). *Pediatr Crit Care Med*. 2007 Sep;8(5):482-5.
257. Finfer S, Chittock DR, Su SY, Blair D, Foster D, Dhingra V; NICE-SUGAR Study Investigators. Intensive versus conventional glucose control in critically ill patients. *N Engl J Med* 2009; 360:1283-1297.
258. Yung M, Wilkins B, Norton L, Slater A, for the Paediatric Study Group, Australian and New Zealand Intensive Care Society. Glucose control, organ failure, and mortality in pediatric intensive care. (*Pediatr Crit Care Med* 2008; 9:147–152).
259. Criado Molina. Diabetes mellitus: manejo y seguimiento en Gascon. Manual práctico de pediatría. Ed. Ergón Madrid. 2008.
260. Yildizdas D, Onenli-Mungan N, Yapicioglu H, Yopaloglu AK, Serdemir Y, Yuskel B. Thyroid hormone levels and their relationship to survival in children with bacterial sepsis and septic shock. *J Pediatr Endocrinol Metab* 2004; 17: 1435-1442.
261. Den Brinker M, Dumas B, Visser TJ, Hop WC, Hazelzet JA, Festen DA. Thyroid function and outcome in children who survived meningococcal septic shock. *Intensive Care Med* 2005; 31: 970-976.
262. Callister ME, Burke-Gaffney A, Quinlan GJ, Betts H, Nadel S, Evans TW. Persistently low plasma thioredoxin is associated with meningococcal septic shock in children. *Intensive Care Med*. 2007; 33(2):364-7.
263. López-Herce J, Bustinza A, Alcazar A, Brandstrup KB, Fernández C y Roncero M. Tratamiento del shock séptico con plasmafiltración y hemodiafiltración continuas simultáneas. *An Pediatr (Barc)* 2003;59(5):491-6.
264. Bock KR. Renal replacement therapy in pediatric critical care medicine. *Curr Opin Pediatr*. 2005 Jun;17(3):368-71.
265. Sociedad Española de Transfusión Sanguínea. Guía sobre la transfusión de componentes sanguíneos y derivados plasmáticos. Sociedad Española de Transfusión Sanguínea. 3ª ed. 2006.

266. Levi M, Toh CH, Thachil J, Watson HG; British Committee for Standards in Haematology. Guidelines for the diagnosis and management of disseminated intravascular coagulation. *Br J Haematol.* 2009 Apr; 145(1):24-33
267. Afshari A, Wetterlslev J, Brok J, Moller AM. Antithrombin III for critically ill patients. *Cochrane database Syst Rev.* 2008 16(3): CD005370.
268. Martí-Carvajal A, Salanti G, Cardona AF. Human recombinant activated protein C for severe sepsis. *Cochrane Database Syst Rev.* 2007 Jul 18;(3): CD004388.
269. Nadel S, Goldstein B, Williams MD, Dalton H, Peters M, Macias WL, Abd-Allah SA, Levy H, Angle R, Wang D, Sundin DP, Giroir B; REsearching severe Sepsis and Organ dysfunction in children: a gLocal perspective (RESOLVE) study group. Drotrecogin alfa (activated) in children with severe sepsis: a multicentre phase III randomised controlled trial. *Lancet.* 2007 Mar 10;369(9564):836-43.
270. Kreymann KG, de Heer G, Nierhaus A, Kluge S. Use of polyclonal immunoglobulins as adjunctive therapy for sepsis or septic shock. *Critical Care Medicine:* 2007; 35:2677-2685.
271. CGIU. Clinical guidelines for Immunoglobulin Use Second Edition. IVIg Guideline Development Group of the IVIg Expert Working Group. 2008: 62-64.
272. Sachs U, Reiter A, Walter T, Bein T, Woessmann W. Safety and efficacy of therapeutic early onset granulocyte transfusions in pediatric patients with neutropenia and severe infections. *Transfusion.* 2006; 46: 1909-1914.
273. Seidel MG, Minkov M, Witt V, Matthes-Martin S, Pötschger U, Worrel N et al. Granulocyte transfusions in children and young adults: does the dose matter? *J Pediatr Hematol Oncol.* 2009; 31(3):166-172.
274. van de Wetering M D, Weggelaar N, Offringa M, Caron H N, Kuijpers T W. Granulocyte transfusions in neutropenic children: a systematic review of the literature. *European Journal of Cancer* 2007; 43(14): 2082-2092
275. Joffe A, Anton N, Lequier L, Vandermeer B, Tjosvold L, Larsen B et al. Nutritional support for critically ill children. *Cochrane Database of Systematic Reviews* 2009, Issue 2. Art. No.: CD005144. DOI: 10.1002/14651858.CD005144.pub2. The Cochrane Library (ISSN 1464-780X).
276. Briassoulis G, Filippou O, Kanariou M, Hatzis T. Comparative effects of early randomized immune or non-immune-enhancing enteral nutrition on cytokine production in children with septic shock. *Intensive Care Med.* 2005;31:851-8
277. Sudarsanam T, Rupali P, Tharyan P, Abraham OC, Thomas K. Administración de antibióticos antes del ingreso para los casos en los que se sospecha una enfermedad meningocócica (Revisión Cochrane traducida). En: *La Biblioteca Cochrane Plus*, 2007 Número 4. Oxford: Update Software Ltd. Disponible en: <http://www.update-software.com>
278. Barberán J, Mensa J, Fariñas C, Linares P, Olaechea P, Palomar M et al. Recomendaciones de tratamiento antimicrobiano en pacientes alérgicos a antibióticos betalactámicos. *Rev Esp Quimioter* 2008;21(1):60-82.
279. Rodrigo C. Sepsis, Sepsis grave y shock séptico. En *Aristegui J Guía de terapéutica antimicrobiana en pediatría 2007-2008. 2ª ed.* Sabadell (Barcelona). Ed Antares 2007. p. 188-191.
280. Ferraris VA. Exploratory laparotomy for potential abdominal sepsis in patients with multiple organ failure. *Arch Surg* 118:1130-1133.
281. Marshall JC, Maier RV, Jimenez M, Dellinger EP. Source control in the management of severe sepsis and septic shock: An evidence-based review. *Crit Care Med.* 2004;32(11 SUPL.):S513-S526.

proceso
asistencial
integrado

