

## 6. BIBLIOGRAFÍA

- Aguilera E.** 1990. Sexual differences in nest attendance and chick-feeding rhythms of white spoonbills. Estación Biológica de Doñana. AUK 107 (2): 416-420.
- Aguilera E., C. Ramo y C. deleCourt** 1996. Food and feeding sites of the Eurasian spoonbill (*Platalea leucorodia*) in southwestern Spain. Colonial Waterbirds 19: 159-166.
- Andral B., J. Y. Stanisiere, D. Sauzade, E. Damier, H. Thebault, F. Galgani, P. Boissery** 2004. Monitoring chemical contamination levels in the Mediterranean based on the use of mussel caging. Marine Pollution Bulletin 49: 704-712.
- Arias A. y P. Drake** 1994. Structure and production of the benthic macroinvertebrate community in a shallow lagoon in the bay of Cádiz. Marine Ecology-Progress Series 115 (1-2): 151-167.
- Arias A. Y Drake P.** 1999. Fauna acuática de las salinas del Parque Natural Bahía de Cádiz. Consejería de Medio ambiente, Junta de Andalucía. 255 pp.
- Arroyo G. M., Hortas F., Pérez-Hurtado A. y De Le Court.** 1997. Cádiz Bay, a new breeding site for Spoonbills (*Platalea leucorodia*) in South Spain: strategies for conservation. Workshop of the Spoonbill network. Eurosite y Societe pour l'Etude et la Protection de la Nature en Bretagne (SEPNB). Sené (Francia).
- Arroyo G. M., Hortas F., de le Court C., Urbina E., Oliveros F., Sánchez I. y Barragán M. A.** 1998. Nueva colonia de cría de la Espátula *Platalea leucorodia* en el Parque Natural de la Bahía de Cádiz. Ardeola 45(1): 119.
- Beyer W. N., M. Spalding y D. Morrison** 1997. Mercury concentrations in feathers of wading birds from Florida. Ambio 26: 97-100.
- Burger J.** 1993. Metals in Avian Feathers: Bioindicators of Environmental Pollution. Rev. Environ.Toxicol. 5: 197-306.
- Burger J.** 1994a. Metals in avian feathers: bioindicators of environmental pollution. Rev. Environ.Toxicol. 5: 302-411.
- Burger J.** 1994b. Heavy metals in avian eggshells: another excretion method. J. Toxicol. Environ.Health 41: 207-220.
- Burger J.** 1996. Heavy metal and selenium levels in feathers of Franklin's Gulls in interior North America. Auk 113: 399-407.
- Burger J.** 1997. Heavy metals and selenium in Herring Gulls (*Larus argentatus*) nesting in colonies from eastern Long Island to Virginia. Environmental Monitoring and Assessment 48: 285-296.

**Burger J.** 2002. Food Chain Differences Affect Heavy Metals in Bird Eggs in Barnegat Bay, New Jersey. *Environmental Research Section A* 90: 33-39.

**Burger J. y M. Gochfeld** 1997. Heavy Metal and Selenium Concentrations in Feathers of Egrets from Bali and Sulawesi, Indonesia. *Arch. Environ. Contam. Toxicol.* 32: 217-211.

**Burger J., G. E. Woolfenden y M. Gochfeld** 1999. Metal Concentrations in the Eggs of Endangered Florida Scrub-Jays from Central Florida. *Arch. Environ. Contam. Toxicol.* 37: 385–388.

**Burger J., R. Bowman, G. E. Woolfenden y M. Gochfeld** 2004. Metal and metalloid concentrations in the eggs of threatened Florida scrub-jays in suburban habitat from south-central Florida. *Science of the Total Environment* 328: 185–193.

**Burger J., T. Shukla, C. Dixon, S. Shukla, M. J. McMahon, R. Ramos y M. Gochfeld** 2001. Metals in feathers of Sooty Tern, White Tern, Gray-backed Tern, and Brown Noddy from islands in the North Pacific. *Environmental Monitoring and Assessment* 71: 71–89.

**Castro G., N. Stoyan y J.P. Myers** 1989. Assimilation efficiency in birds - a function of taxon or food type. Comparative biochemistry and physiology a-physiology 92 (3): 271-278.

**Castro M.** 2001. Ecología de la alimentación en el Chorlitejo patinegro *Charadrius alexandrinus*. ¿Existe plasticidad comportamental y fisiológica asociada al sexo?. Tesis Doctoral. Universidad de Cádiz.

**David P.G.** 1994. Wading bird use of lake Okeechobee relative to fluctuating water levels. *Wilson Bull* 106(4): 719-732.

**De la Court C.** 2004. Spoonbills in Spain in 2004. Spoonbill Network Eurosite. Newsletter nº 2.

**Del Valls T. A., J. M. Forja, E. González-Mazo, A. Gómez-Parra y J. Blasco** 1998. Determining contamination sources in marine sediments using multivariate analysis. *Trends in analytical chemistry*. 17: 181-192.

**Del Valls T. A., J.M. Forja y A. Gómez-Parra** 1998. Integrative assessment of sediment quality in two littoral ecosystems from the Gulf of Cádiz, Spain. *Environmental Toxicology and Chemistry*. 17: 1073–1084.

**Drake P. y A. Arias** 1995. Distribution and production of *Chironomus salinarius* (diptera, chironomidae) in a shallow coastal lagoon in the bay of Cádiz. *Hydrobiologia* 299 (3): 195-206.

**Durell S.E.A.LD., S. McGrorty, A.D. West, R.T. Clarke, J.D. Goss-Custard y R.A. Stillman** 2005. A strategy for baseline monitoring of estuary Special Protection

Areas. Biological Conservation 121 (2): 289-301.

**Eisler R.** 1987. Mercury Hazards to Fish, Wildlife, y Invertebrates: A Synoptic Review. U.S. Fish and Wildlife Service. *Biolog. Rep.* 85 (1.1).

**Eisler R.** 2000. Handbook of chemical risk assessment: health hazards to humans, plants and animals. Boca Raton, FL: Lewis Publishers.

**Eldridge J.** 1992. Management of habitat for breeding and migrating shorebirds in the midwest. University of Minnesota. Fish and Wildlife Leaflet. 13.2.14.

**Equipo de Seguimiento de Acuáticas de Cádiz** 2004. Seguimiento de la colonia de reproducción de la Espátula Común *Platalea leucorodia* en la Isla del Trocadero (Salina La Covacha).

**Equipo de Seguimiento de Acuáticas de Cádiz** 2005. Propuesta complementaria al plan andaluz de acuáticas para la realización de censos. Consejería de Medio Ambiente. Informe censo de invernantes. Bahía de Cádiz.

**Equipo de Seguimiento de la Estación Biología del Doñana** 2003. Anillamiento de la Espátula Común *Platalea leucorodia* en el P.N. de la Bahía de Cádiz (La Covacha).

**Estrella S.M.** 2001. Comportamiento alimentario, selección de presa y balance energético de la Aguja colinegra *Limosa limosa* en hábitats supramareales de la Bahía de Cádiz. Tesis de licenciatura. Departamento de Biología. Universidad de Cádiz.

**Franco A.R. y Rodriguez M.** 2001. Libro Rojo de los vertebrados amenazados de Andalucía. Consejería de Medio Ambiente de la Junta de Andalucía.

**Furness R. W., S.J. Muirhead y M. Woodburn** 1986. Using bird feathers to measure mercury in the environment: relationships between mercury content and moult. *Mar Pollut Bull* 17: 27-30.

**Glassom D. y G.M. Branch** 1997. Impact of predation by greater flamingos *Phoenicopterus ruber* on the meiofauna, microflora, and sediment properties of two southern African lagoons. *Marine Ecology Progress Series* 150: 1-10.

**Gochfeld M., J.I. Belan, T. Shukla, T. Benson y Burger J.** 1996. Heavy metals in laughing gulls: gender, age and tissue differences. *Environmental Toxicology and Chemistry* 15: 2275-2283.

**Gochfeld M. y J. Burger** 1998. Temporal Trends in Metal Levels in Eggs of the Endangered Roseate Tern (*Sterna dougallii*) in New York. *Environmental Research, Section A* 77: 36-42.

**Goss-Custard J.D., R.T. Clarke, K.B. Briggs, B.J. Ens, K.M. Exo, C. Smit, A.J. Beintema, R.W.G. Caldow, D.C. Catt, N.A. Clark, S. Durell, M.P. Harris, J.B.**

**Hulscher, P.L. Meininger, N. Picozzi, R. Prysjones, U.N. Safriel, A.D. West** 1995. Population consequences of winter habitat loss in a migratory shorebird .1. Estimating model parameters. *Journal of applied ecology* 32 (2): 320-336.

**Goss-Custard J.D., R.A. Stillman, R.W.G Caldow, A.D. West y M. Guillemain** 2003. Carrying capacity in overwintering birds: when are spatial models needed?. *Journal of Applied Ecology* 40 (1): 176-187.

**Goutner V. y R.W. Furness** 1997. Mercury in Feathers of Little Egret *Egretta garzetta* and Night Heron *Nycticorax nycticorax* Chicks and in Their Prey in the Axios Delta, Greece. *Arch. Environ. Contam. Toxicol.* 32: 211–216.

**Grupo de Conservación de Humedales Costeros (GEAM)** 1996. Cartografiado de las aves invernantes y reproductoras en el Parque Natural de la Bahía de Cádiz. Informe inédito. Universidad de Cádiz-Consejería de Medio Ambiente de la Junta de Andalucía.

**Grupo de Conservación de Humedales Costeros (GEAM)** 1998. Incidencias de la Actividad Marisquera sobre las Poblaciones de Aves e Invertebrados Bentónicos Intermareales en el Parque Natural Bahía de Cádiz. Informe inédito. Universidad de Cádiz-Consejería de Medio Ambiente de la Junta de Andalucía.

**Grupo de Conservación de Humedales Costeros (GEAM)** 1999. inventario de aves del Parque Natural Bahía de Cádiz. Informe Inédito. Proyecto Life-Nature "Oiesaux d'eau de la façade atlantique". Universidad de Cádiz-Consejería de Medio Ambiente de la Junta de Andalucía.

**Grupo de Conservación de Humedales Costeros (GEAM)** 2000. Maintaining migratory coastal bird diversity: management through behaviour-based predictive modelling. Comisión Europea. V Programa Marco Programa "Energy, Environment and Sustainable Development". Centre for Ecologyand Hydrology(Reino Unido). Universidad de Cádiz (España),Groupe d'Etude del Milieux Estuarinens et Littoraux (Francia), Alterra (Holanda), National Environment Research Institute (Dinamarca). University of Reading (Reino Unido) y Centre National de la Recherche Scientifique (Francia).

**Grupo de Conservación de Humedales Costeros** 2003. Cartografiado de las aves invernantes y reproductoras en el Parque Natural Bahía de Cádiz. Año 2002. Informe inédito. Universidad de Cádiz y Consejería de Medio Ambiente de la Junta de Andalucía.

**Grupo de Conservación de Humedales Costeros** 2004a. Plan experimental de recuperación, manejo y usos sostenibles de humedales costeros marismeños en el Parque Natural Bahía de Cádiz. Universidad de Cádiz y Consejería de Medio Ambiente de la Junta de Andalucía.

**Grupo de Conservación de Humedales Costeros** 2004b. Maintaining migratory coastal bird diversity: Management though behaviour-based predictive

population modelling. Universidad de Cádiz. Proyecto financiado por el V Programa Marco de la Unión Europea.

**Helmers D.L.** 1991. Habitat use by migrant shorebirds and invertebrate availability in a managed wetland complex. M.S. Thesis, Univ. of Missouri Columbia. 135pp.

**Helmers D.L.** 1992. Shorebird management manual. Western Hemisphere Shorebird Reserve Network, Manomet, MA. 58 pp.

**Hernández L. M., B. Gómara, M. Fernández, B. Jiménez, M. J. González, R. Baos, F. Hiraldo, M. Ferrer, V. Benito, M. A. Suñer, V. Devesa, O. Muñoz y R. Montoso** 1999. Accumulation of heavy metals and As in wetland birds in the area around Doñana National Park affected by the Aznalcollar toxic spill. *The Science of the Total Environment* 242: 293-308.

**Hortas F.** 1997. Evolución de la comunidad de aves limícolas (Orden Charadriiformes) en salinas del suroeste de España. Estructura espacio-temporal de las poblaciones y uso del hábitat. Tesis Doctoral. Universidad de Cádiz.

**Hortas F. y Arroyo G. M.** 2000. Status and management plan of Spoonbill (*Platalea leucorodia*) populations in Cádiz, Southwest Spain. 3th Eurosite Spoonbill Workshop "Wetlands management and conservation of Spoonbills and other waterbirds". La Rábida, Abril 2000. Huelva.

**Jong-Ryol C., I. Tsuchiya y H. Sugita** 2000. Captive breeding of Black-faced Spoonbills. Pages 47-53 in Conservation and research of Black-faced Spoonbills and their habitats, second edition. (M. Ueta, R. Kurosawa y D. Allen, Eds.). Wild Bird Society of Japan, Tokyo, Japan.

**Karasov W.H.** 1990. Digestion in birds: chemical and physiological determinants and ecological implications en *Avian foraging: theory, methodology and applications* (Eds. M.L. Morrison, C.J. Ralph, J.Verner y J.R. Jehl) Studies in Avian Biology nº 13:391-415. Cooper Ornithological Society, Lawrence, Kansas.

**Lewis S. A. y R. W. Furness** 1991. Mercury accumulation and excretion in laboratory reared black-headed gull *Larus ridibundus* chicks. *Arch Environ Contam Toxicol.* 21: 316-320.

**Masero J.A.** 2002. Why don't Red Knots *Calidris canutus* feed extensively on the crustacean *Artemia*? . *Bird Study* 49: 304-306.

**Masero J.A.** 2003. Assessing alternative anthropogenic habitats for conserving waterbirds: salinas as buffer areas against the impact of natural habitat loss for shorebirds. *Biodiversity and conservation* 12 (6): 1157-1173.

**Masero J.A., M. Pérez-González, M. Basadre, y M. Otero-Saavedra** 1999. Food supply for waders (Aves: Charadrii) in an estuarine area of the Bay of Cádiz (SW

Iberian Peninsula). *Acta Oecologica*. 20:429-434.

**Masero J.A, A. Pérez-Hurtado, M. Castro y G. Arroyo** 2000. Complementary use of intertidal mudflats and adjacent salinas by foraging waders. *Ardea* 88:177-191.

**Masero J.A. y A. Pérez-Hurtado** 2001. Importance of the supratidal habitats for maintaining overwintering shorebird populations: how redshanks use tidal mudflats and adjacent saltworks in southern Europe. *Condor* 103: 21-30.

**Matheu E. y J. del Hoyo** 1992. Family Threskiornithidae (Ibis and Spoonbills). Páginas 472-506 en *Handbook of the Birds of the World*, vol 3 (J. Del Hoyo., A. Elliott, y J. Sargatal, Eds.). Hoatzin to Auks. Lynx Editions, Barcelona.

**McArthur R.H. y E.R. Pianka** 1966. On optimal use of a patchy environment. *American Naturalist* 100: 603-609.

**Méndez N., Romero J y J. Flos** 1997. Population dynamics of the polychaete *Capitella capitata* in the littoral zone of Barcelona (Spain, NW Mediterranean). *Journal of Experimental Marine Biology and Ecology* 218: 263-284.

**Monteiro L. R. y R.W. Furness** 1997. Accelerated increase in mercury contamination in North Atlantic mesopelagic food chains as indicated by time series of seabird feathers. *Environmental Toxicology and Chemistry* 16: 2489-2493.

**Palma L., P. Beja, P.C. Tavares, y L.R. Monteiro** 2005. Spatial variation of mercury levels in nesting Bonelli's eagles from Southwest Portugal: effects of diet composition and prey contamination. *Environmental Pollution* 134 : 549-557.

**Pérez-Hurtado A., J. A. Masero, G.M. Arroyo, S. Rodríguez, F. Hortas y M. Castro** 1997. Estudio de la incidencia de la avifauna ictiófaga en los cultivos marinos y otras especies orníticas en el Parque Natural bahía de Cádiz. Consejería de Medio Ambiente de la Junta de Andalucía.

**Pérez-Hurtado A., J.A. Masero, G.M. Arroyo, F. Hortas y M. Castro** 1997. Importancia de las poblaciones de aves limícolas en el Parque Natural Bahía de Cádiz. Bases para su declaración como área Ramsar. En la ruta migratoria del Atlántico Oriental. Eds. Amigos de la Tierra y Junta de Andalucía pp: 185-192. Madrid.

**Peter G. David** 1994. Wading bird use of lake Okeechobee relative to fluctuating water levels. *Wilson Bull* 106(4): 719-732.

**Piersma T., de Goeij P. y I. Tulp** 1993. An evaluation of intertidal feeding habitats from a shorebird perspective: towards relevant comparisons between temperate and tropical mudflats. *Netherlands Journal of Sea Research* 31: 503-512.

**Ricklefs R.E.** 1967. A graphical method of fitting equations to growth curves. *Ecology*, 48: 978.

**Ricklefs R.E.** 1968. Patterns of growth in birds. *Ibis*, 110: 419-451.

**Rosenberg R., M. Blomqvist, H. C. Nilsson, H. Cederwall y A. Dimming** 2004. Marine quality assessment by use of benthic species-abundance distributions: a proposed new protocol within the European Union Water Framework Directive. *Marine Pollution Bulletin* 49: 728-739.

**Ruiz F. J., Almorza D. y Fernández Zapata, J.** 1990. The salt-pans of the Bahía de Cádiz as a new and unique nesting habitat for Herring Gulls *Larus argentatus*. *Alectoris* 7: 70-76.

**Sadoul N., Walmsley J. y Champentier B.** 1998. Salinas and Nature Conservation. *Conservation of Mediterranean Wetlands*, number 9. MedWet/Tour du Valat Publications, Arles, France.

**Shannon C. E. y Weaver W.** 1949. The mathematical theory of communication. Illinois Press., Urbana.

**Sokolowski A., M. Wolowicz, H. Hummel, K. Smolarz-Górska, D. Fichet, G. Radenac, C. Thiriot-Quiévreux, J. Namiésnik** 2004. Abnormal features of *Macoma balthica* (*Bivalvia*) in the Baltic Sea: alerting symptoms of environmental adversity?. *MARINE POLLUTION BULLETIN* 49: 17-22.

**Stephen D.W. y J.R. Krebs** 1989. Foraging theory. Princeton University Press, Princeton, New Jersey.

**Swennen C. y Y-T. Yu** 2005. Food and feeding behaviour of the Black-faced Spoonbill. *Waterbirds* 28: 19-27.

**Tavares P. C., L. R. Monteiro, R. J. Lopes, M. M. C. Santos y R. W. Furness** 2004. Intraspecific variation of mercury contamination in chicks of Black-winged Stilt (*Himantopus himantopus*) in coastal wetlands from Southwestern Europe. *Bulletin of Environmental Contamination and Toxicology* 72(3): 437-444.

**Tavares P.C. , L.R. Monteiro, R.J. Lopes, M.E. Pereira, A.C. Duarte y R.W. Furness** 2005. Variation of mercury contamination in chicks of Little Tern *Sterna albifrons* in southwest Europe : brood, age, and colony related effects. *Bulletin of Environmental Contamination and Toxicology* 74(1): 177-183.

**Thompson D. R., R.W. Furness y L.R. Monteiro** 1998. Seabirds as biomonitor of mercury inputs to epipelagic and mesopelagic marine food chains. *The Science of the Total Environment* 213: 299-305.

**Thyren S., P.H. Becker y H. Behmann** 2000. Organochlorine and mercury contamination of little terns (*Sterna albifrons*) breeding at the western Baltic Sea, 1978-96. *Environmental Pollution* 108: 225-238.

**Tori G. M., S. McLeod, K. McKnight, T. Moorman, F.A. Reid** 2002. Wetland Conservation and ducks unlimited: Real world approaches to multispecies management. *Waterbirds (Waterbirds)* 25(2): 115-121.

**Van Gils J.A., I.W. Schenk, O. Bos y T. Piersma** 2003. Incompletely informed shorebirds that face a digestive constraint maximize net energy gain when exploiting patches. *American Naturalist* 161 (5): 777-793.

**Wader Study Group** 2003. Are waders world-wide in decline? Reviewing the evidence. Conclusions of the International WSG Annual General Meeting, Cádiz 25-28 Septiembre. *Wader Study Group Bulletin* 101-102: 2-13.

**Warwick R.M** 2001. Evidence for the effects of metal contamination on the intertidal macrobenthic assemblages of the Fal Estuary. *Marine Pollution Bulletin* 42: 145-148.

**West A.D., J.D. Goss-Custard, S. Durell y R.A. Stillman** 2005. Maintaining estuary quality for shorebirds: towards simple guidelines. Centre for Ecology and Hydrology for Dorset, Winfrith Newburgh, Dorchester, UK. *Biological Conservation* 123: 211-224.

**Yu Y. T. y C. Swennen** 2004. Feeding of wintering Black-faced Spoonbills in Hong Kong: when and how long? *Waterbirds* 27: 135-140.

**Zwarts L. y S. Dirksen** 1990. Digestive Bottleneck Limits the Increase in Food-intake of Whimbrels Preparing for Spring Migration from the Banc-Darguin, Mauritania. *ARDEA* 78 (1-2): 257-278.